

Accepted Manuscript


Title: A systematic review of the neurophysiology of mindfulness on EEG oscillations

Author: Tim Lomas Itai Ivztan Cynthia H.Y. Fu

PII: S0149-7634(15)00251-1

DOI: <http://dx.doi.org/doi:10.1016/j.neubiorev.2015.09.018>

Reference: NBR 2278



To appear in:

Received date: 15-5-2015

Revised date: 23-9-2015

Accepted date: 30-9-2015

Please cite this article as: Lomas, T., Ivztan, I., Fu, C.H.Y., A systematic review of the neurophysiology of mindfulness on EEG oscillations, *Neuroscience and Biobehavioral Reviews* (2015), <http://dx.doi.org/10.1016/j.neubiorev.2015.09.018>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

- We conducted a systematic review of EEG studies of mindfulness meditation
- We examined power differentials between mindfulness and a control state
- Mindfulness was associated with enhanced alpha and theta power
- No consistent patterns were observed in terms of beta, delta and gamma
- Elevated alpha and theta may signify a state of relaxed alertness

Accepted Manuscript

Running title: The neurophysiology of mindfulness

Title

1 A systematic review of the neurophysiology of mindfulness on EEG oscillations
2
3
4
5

Authors

6
7
8 Tim Lomas ¹, Itai Ivtzan ¹, Cynthia H.Y. Fu ^{1,2}
9

Affiliations:

- 10
11
12
13
14
15 1. School of Psychology, University of East London, London, UK
16
17 2. Centre for Affective Disorders, Institute of Psychiatry, Psychology and Neuroscience,
18
19 King's College London, London, UK
20
21
22
23

Author for Correspondence:

24
25 Dr. Tim Lomas, School of Psychology, University of East London, Arthur Edwards Building,
26
27 Water Lane, London, E15 4LZ, United Kingdom
28

29 Tel: +44 (0)20 8223 4465 Fax: +44 (0)20 8223 4937 Email: t.lomas@uel.ac.uk
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Running title: The neurophysiology of mindfulness

Abstract:

1
2
3 Mindfulness meditation has been purported as a beneficial practice for wellbeing. It would
4 be expected that the neurophysiology of mindfulness would reflect this impact on wellbeing.
5
6 However, investigations of the effect of mindfulness have generated mixed reports of
7
8 increases, decreases, as well as no differences in EEG oscillations in comparison with a
9
10 resting state and a variety of tasks. We have performed systematic review of EEG studies of
11
12 mindfulness meditation in order to determine any common effects and to identify factors
13
14 which may impact on the effects. Databases were reviewed from 1966 to August 2015.
15
16 Eligibility criteria included empirical quantitative analyses of mindfulness meditation practice
17
18 and EEG measurements acquired in relation to practice. A total of 56 papers met the
19
20 eligibility criteria and were included in the systematic review, consisting of a total 1,715
21
22 subjects: 1,358 healthy individuals and 357 individuals with psychiatric diagnoses. Studies
23
24 were principally examined for power outcomes in each bandwidth, in particular the power
25
26 differentials between mindfulness and the control state, as well as outcomes relating to
27
28 hemispheric asymmetry and event-related potentials. The systematic review revealed that
29
30 mindfulness was most commonly associated with enhanced alpha and theta power as
31
32 compared to an eyes closed resting state, although such outcomes were not uniformly
33
34 reported. No consistent patterns were observed with respect to beta, delta and gamma
35
36 bandwidths. In summary, mindfulness is associated with increased alpha and theta power in
37
38 both healthy individuals and in patient groups. This co-presence of elevated alpha and theta
39
40 may signify a state of relaxed alertness which is conducive to mental health.
41
42
43
44
45
46
47

Keywords: mindfulness; meditation; neurophysiology; EEG; systematic review.

Running title: The neurophysiology of mindfulness

Introduction

1
2
3 Meditation refers to a diverse range of mental activities which share a common focus on the
4 regulation of attention and awareness (Cahn and Polich, 2006) in order to improve voluntary
5 control of mental processes which is purported to foster general wellbeing (Walsh and
6 Shapiro, 2006). Most world cultures have developed their own forms of meditation; for
7 example, Christianity has a long tradition of contemplative prayer (Egan, 1978). Much of the
8 recent scientific interest in meditation has centred on mindfulness meditation, which is a
9 practice that is believed to have originated with Buddhism around the fifth millennium B.C.
10 although its roots may stretch back further to the third millennium B.C. in Hindu culture
11 (Cousins, 1996).
12
13
14
15
16
17
18
19
20
21
22
23

24 The most common forms of meditation may be conceptualized as involving either focused
25 attention or an open-monitoring form of processes (Lutz et al., 2008). Focused attention
26 practices may be operationalized into their respective attention networks (Posner and
27 Petersen, 1990; Mirsky et al., 1991): sustained attention (e.g. towards a target, such as the
28 breath), executive attention (e.g. preventing one's focus from 'wandering'), attention
29 switching (e.g. disengaging from distractions), selective attention and attention re-orienting
30 (e.g. redirecting focus back to the breath), and working memory (Lutz et al., 2008; Vago and
31 Silbersweig, 2012). Open-monitoring refers to a broader receptive awareness or capacity to
32 detect events within an unrestricted awareness without a specific focus (Raffone and
33 Srinivasan, 2010), which can include a process of 'meta-awareness' (i.e., awareness of
34 awareness, in which practitioners are able to reflect on the process of consciousness itself).
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50

51 Mindfulness has been described as the awareness that arises through purposeful attention
52 on the present moment with nonjudgmental experience (Kabat-Zinn, 2003). While
53 mindfulness has been commonly viewed as an example of open-monitoring, it has been
54 proposed to involve an admixture of focused attention and open-monitoring (Lutz et al.,
55 2008; Vago and Silbersweig, 2012) as most mindfulness practices begin with a period of
56
57
58
59
60
61
62
63
64
65

Running title: The neurophysiology of mindfulness

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

focused attention on a target, such as the breath, in order to focus awareness, followed by the more receptive state of open-monitoring (Cahn and Polich, 2006). In Vago and Silbersweig's model (2012), the practice of mindfulness leads to three overarching self-related capacities: meta self-awareness, self-regulation, and self-transcendence. These are subserved by numerous subcomponent cognitive components, including motivation (which is crucial in terms of people practicing meditation in the first place), attention regulation (via the development of attention modalities), and de-centring (an ability, defined below, that arises from enhanced attention regulation, and which facilitates self-awareness and transcendence). It is further proposed that these three overarching capacities modulate 'self-specifying and narrative self-networks' through an integrative fronto-parietal control network.

Mindfulness has been applied as a clinical intervention based on the notion that it is a method for training attention and awareness. By developing the ability to observe one's thoughts and feelings, practitioners learn how to perceive them as temporary, objective events in the mind as opposed to reflections of the self that are necessarily true, which has been termed as the ability to "decentre" (Fresco et al., 2007). As a clinical intervention, it involves the process to engage with negative experiences, such as pain or dysphoric emotions, with more dispassion and less reactivity (Shapiro et al., 2005). Mindfulness was initially applied as an intervention for chronic pain with Kabat-Zinn's (1982) Mindfulness-Based Stress Reduction (MBSR) program. The MBSR program has since been applied in the treatment for number of conditions, including cancer (Ledesma & Kumano, 2009) and migraine (Schmidt et al., 2010), and adapted as a treatment to prevent relapse in depression (Mindfulness-Based Cognitive Therapy; Segal et al., 2002) and for the treatment of substance abuse (Mindfulness-Based Relapse Prevention; Bowen et al., 2014, Mindfulness-Oriented Recovery Enhancement; Garland et al., 2014).

The effectiveness of mindfulness has been assessed by measures for depression and quality of life (Hofmann et al., 2010). As mindfulness may be considered to be a method of

Running title: The neurophysiology of mindfulness

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

attention training and emotion regulation, we would expect that the corresponding neurophysiological states should be observable. Electroencephalography (EEG) is a non-invasive technique that analyzes spatiotemporal aspects of underlying brain activity, which provides a measure of the large-scale synchronization of neural networks (Cacioppo et al., 2007). Patterns of EEG activity to particular meditative states have been investigated. A commonly reported feature of meditation has been theta and alpha event-related synchronization (Fell et al., 2010), which are regarded as markers of internally-directed attention processing (Shaw, 1996). Such synchronization has been observed across different meditation practices, including mindfulness, as well as practices such as transcendental meditation, which involves focused attention upon an internally-voiced mantra. However, different types of meditation practice have been associated with unique frequency patterns, reflecting the form of attention (Dunn et al., 1999). For example, mindfulness has been associated with increase alpha power while focused attention has been associated with increased gamma activity and idiosyncratic meditation with decreased alpha and beta (Hinterberger et al., 2014).

Additionally, Event-Related Potentials (ERP) provide a measure of large number of time-locked experimental trials, enabling the analysis of sensory, perceptual, and cognitive processing (Light et al., 2010). Such studies involve the precision analysis of populations of neuronal transients directly manifested via a stimulus/event, which is frequently a stimulus connected to an attention-based task (e.g., listening to an auditory signal) (Schoenberg and Speckens, 2014). The high temporal resolution of this approach, involving millisecond precision, allows the investigation of early information processing stages and subsequent transitions to higher-level cognitive operations. ERP studies have been used to corroborate the idea of mindfulness as a system of attention training. For example, van Leeuwen et al. (2012) examined the impact of mindfulness practice on hierarchical stimulus processing and attentional selection, focusing on differences in early components of the evoked visual response (e.g., P1 and N1 components) in meditators versus matched controls. Meditators

Running title: The neurophysiology of mindfulness

1 exhibited faster attentional disengagement from a dominant global presentation in order to
2 focus in on specific stimuli, suggesting that meditation enhances speed of attention
3 allocation and relocation, thus increasing the depth of information processing.
4
5

6
7 In the present review, we have focused on mindfulness meditation. We have examined
8 factors which appear to impact upon EEG measures including the experience of the
9 meditator, being a novice or relative expert, as experience has been reported to accentuate
10 amplitude differences between meditation and the resting state (Hinterberger et al., 2014)
11 while the converse has also been observed (Cahn et al., 2010). An additional factor
12 includes the location of the brain activity. For example, increased alpha during mindfulness
13 has been localized to frontal regions (Takahashi et al., 2005) but has also been observed
14 increases in posterior regions (Lagopoulos et al., 2009; Cahn et al., 2010). Furthermore,
15 EEG analysis of meditation may be affected by whether the control task is a resting state or
16 a cognitive task as increased theta amplitude during meditation has been observed in
17 comparison to a resting state baseline, but was comparable in amplitude to an executive
18 attention task, which may be further modulated by the experience of the meditator (Lomas et
19 al., 2014).
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37

38 We sought to perform a systematic review of patterns of electrophysiological activity
39 associated mindfulness in order to examine the impact on neurophysiology as assessed by
40 EEG bandwidth activation and other measures, including hemispheric asymmetry or event-
41 related potential, and the functional significance of these activities. If mindfulness is
42 expected to impact on functioning attentional networks as well as open-monitoring, then we
43 would expect to observe distinct neural features associated with its practice. We also
44 expected that the experience of the meditator, type of control task, and location of the EEG
45 oscillation would moderate the impact of mindfulness on neurophysiology.
46
47
48
49
50
51
52
53
54
55
56
57
58
59

60 **Methods**

Running title: The neurophysiology of mindfulness

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

The literature search was conducted using the MEDLINE and Scopus electronic databases with the criteria: “EEG” (AND) “mindfulness OR meditation”, in all fields in MEDLINE and limited to article title, abstract, and keywords in Scopus, with the dates: from 1966 to 1st August 2015. The participants, interventions, comparisons, outcomes and study design (PICOS) characteristics, the key criteria were interventions: mindfulness meditation or functional equivalent; participants: adults; and outcomes: EEG analysis. Studies were required to be published, or a manuscript in press, and to be in English. The review was conducted according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines (Moher et al., 2009). The review protocol was registered with the International Prospective Register of Systematic Reviews (PROSPERO) database on 15th September 2014. Registration number: CRD42014013766 (<http://www.crd.york.ac.uk/PROSPERO>).

Inclusion criteria were: 1) mindfulness meditation practice or functional equivalent, such as Vipassana or Zen meditation; 2) EEG measurements acquired in relation to mindfulness meditation practice (whether assessment during the practice itself or connected to its practice, e.g., pre- and post-intervention); 3) quantitative analysis supported by appropriate statistical methodology; and exclusion criteria; and 4) adult sample; and exclusion criteria: 1) theoretical articles or commentaries without statistical analyses.

The following variables were extracted from each paper: experimental protocol (control condition, meditation condition, and/or experimental task), experience of participants (novice or expert), sample features (clinical or non-clinical), outcomes for each individual bandwidth (alpha, beta, theta, delta, and gamma), hemispheric asymmetry, and any event-related potential outcomes.

The primary summary measures were differences in levels of power in each of the bandwidths. Neural activity generates electrical potentials which can be analyzed in terms of parameters of amplitude, frequency, coherence and synchrony. Amplitude, or power, which

Running title: The neurophysiology of mindfulness

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

is the square of the amplitude, reflects the magnitude of the electrical signal, representing the level of synchronized activity in the underlying tissue, i.e. neurons discharging simultaneously. Frequency is the number of oscillatory cycles per second and is divided into the following bandwidths: Delta (1-4 Hz); Theta (4-8 Hz); Alpha (8-13 Hz); Beta (13–30 Hz); and Gamma (36-44 Hz) (Cacioppo et al., 2007). EEG connectivity is the functional integration of spatially distributed neural populations which can be assessed in terms of synchrony, the degree of leading or lagging in the relationship between signals from electrode pairs, and coherence, the stability of that phase relationship.

The primary summary variable was principally the difference in power between a meditation condition and a resting state condition. Secondary power differentials included longitudinal pre- and post- differences, such as, in meditation and/or resting state and/or task conditions before and after an intervention. If applicable, outcomes relating to coherence, synchrony, asymmetry and event-related potentials were also noted.

Of note, there was considerable diversity in how the experience of the participant was defined. In terms of years meditating, the range for which papers rated participants as being 'experienced' varied from 1 year (Kasamatsu and Hirai, 1966) to 9 years (Lagopoulos et al., 2009). Likewise, in terms of hours meditating, the range for which papers rated participants as being 'experienced' varied from 40 hours (Hinterberger et al., 2011) to 1740 hours (Berkovich-Ohana et al., 2012). In the present systematic review, we have applied the lowest of these cutoffs, such that an 'experienced' (i.e., non-novice) meditator was considered to have been meditating for longer than 1 year or have completed more than 40 hours of meditation.

The Quality Assessment Tool for Quantitative Studies (QATQS; National Collaborating Centre for Methods and Tools, 2008) was used to assess the quality of the studies. QATQS assesses methodological rigor in six areas: (a) selection bias; (b) design; (c) confounders; (d) blinding; (e) data collection method; and (f) withdrawals and drop-outs. Each area is

Running title: The neurophysiology of mindfulness

1
2 assessed on a quality score of 1 to 3 (1 = strong; 2 = moderate; 3 = weak). Scores for each
3 area were collated, and a global score was assigned to each study. If there are no weak
4 ratings, the study is given a score of 1 (judged as strong); one weak rating leads to a score
5 of 2 (moderate); and two or more weak ratings generates a score of 3 (weak)
6
7

8 (Supplementary Materials). QATQS scoring was conducted (II) and checked independently
9 (TL). Any discrepancies were resolved by discussion with agreement reached in all cases.
10
11

12
13 The first authors of each paper were contacted for additional information as needed (Amihai
14 and Kozhevnikov, 2014; Arita, 2012; Cahn et al., 2010; Cahn et al., 2013; Hinterberger et al.,
15
16 2011; Hinterberger et al., Walach, 2014; Howells et al., 2012; Huang and Lo, 2009;
17
18 Lagopoulos et al., 2009; Lehmann et al., 2012; Lo et al., 2003; Milz et al., 2014; Murata et
19
20 al., 2004; Saggat et al., 2012; Stinson and Arthur, 2013; Tang et al., 2009; Xue et al., 2014).
21
22

23 Data were extracted (TL) and reviewed (II) with guidance and review (CF).
24
25
26
27
28
29
30

31 **Results**

32 ***Search results***

33
34
35 Following removal of duplicate citations, 284 potentially relevant papers were identified (302
36 articles from Scopus, 291 articles from MEDLINE, and 12 from the reference lists of articles).
37
38

39 From the abstract review, 120 papers were excluded. From the full text reviews of 164
40 papers, 108 papers were excluded. Thus, a total of 56 papers were included in the
41 systematic analysis. Ten of these papers were identified as reporting on overlapping
42 samples: (Berkovich-Ohana et al., 2012; Berkovich-Ohana et al., 2013); (Cahn et al., 2010;
43 Cahn et al., 2013); (Slagter et al., 2007; Slagter et al., 2009); (Hinterberger et al., 2011;
44 Hinterberger et al., 2014); (Schoenberg and Speckens, 2014; Schoenberg and Speckens,
45 2015). As such, the 56 papers included in the systematic analysis represented results from
46 51 independent participant samples ($n = 1,715$ subjects; age range = 19-72 years) (Figure
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

Running title: The neurophysiology of mindfulness

1
2 samples ($n = 1,358$ subjects; age range = 18-72 years)(Table 1), and 10 papers included
3 participants with a psychiatric disorder, representing results from 9 independent samples (n
4 = 357 subjects; age range = 22-64 years): 3 studies on depressed patients in remission ($n =$
5 157), 1 study of patients with suicidal depression ($n = 22$), 1 study involving patients
6 diagnosed with major depressive disorder, reported across 2 papers (Schoenberg and
7 Speckens, 2014, 2015) ($n = 51$), 1 study of patients with bipolar disorder ($n = 21$), 1 study of
8 patients with chronic pain ($n = 27$), 1 study of patients with chronic pain with risk of opioid
9 abuse ($n = 29$), and 1 study of patients with attention-deficit/hyperactivity disorder (ADHD) (n
10 = 50) (Table 2).

11 The findings fall into two main types: (a) studies examining the effects of mindfulness in
12 comparison with a resting state; and (b) studies examining longitudinal changes in EEG
13 patterns relating to practicing mindfulness (Table 3, Supplementary Tables 3-9).

24 ***Effects of mindfulness on neurophysiology***

25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
Twenty-one studies examined the alpha bandwidth, reporting greater amplitude during
mindfulness in comparison with an eyes-closed resting state ($n = 12$), lower amplitude ($n =$
1), and no significant differences ($n = 3$) (Table 3). Most of the studies involved experienced
meditators; novice participants were involved in 4 of the reports of greater amplitude and 1 of
the reports of no significant differences. Coherence was examined in 2 papers, with mixed
results, and more complex analyses in another 2 papers.

The beta bandwidth was examined in 12 studies which compared mindfulness with a resting
state, reporting greater amplitude during mindfulness ($n = 3$; including $n = 1$ with novice
meditators), lower amplitude ($n = 1$), and no significant differences ($n = 5$, including $n = 2$
novices). Coherence ($n = 1$, with no difference found), asynchrony ($n = 1$, finding higher
synchrony with meditation) and more complex analyses ($n = 1$) were also examined.

The theta bandwidth was examined in 19 studies, reporting greater amplitude during
mindfulness ($n = 11$; including $n = 3$ with novices), lower amplitude ($n = 3$; including $n = 2$

Running title: The neurophysiology of mindfulness

1
2 with novices), and no significant differences ($n = 2$; $n = 1$ with novices). Coherence ($n = 2$,
3 with no difference found), and more complex analyses ($n = 1$) were also examined.

4
5 The delta bandwidth was examined in 5 studies, reporting greater amplitude during
6 mindfulness ($n = 1$, with novices, limited to frontal regions) as well as no significant
7 differences ($n = 3$; $n = 1$ with novices). More complex analyses ($n = 1$) were also examined.

8
9
10
11
12 The gamma bandwidth was examined in 7 studies, reporting greater amplitude during
13 mindfulness ($n = 3$) and no significant differences ($n = 2$; $n = 1$ with novices). Gamma
14 amplitude during mindfulness also correlated with train mindfulness and years of practice (n
15 = 1). Coherence ($n = 1$, with no difference found) and asymmetry ($n = 3$, finding greater left-
16 sided activation ($n = 2$) and no differences ($n = 1$)) were also examined.

17
18
19
20
21
22
23
24
25 Event-related potentials were examined in 15 studies, with mindfulness found to have an
26 impact on attention processing measures including P300 ($n = 5$; $n = 2$ on P3b specifically),
27 Late Positive Potential ($n = 2$), Feedback Related Negativity ($n = 1$), Error Related Negativity
28 ($n = 1$), Readiness Potential ($n = 1$), pain-evoked ERPs ($n = 2$), Late Contingent Negative
29 Variation ($n = 1$), and a Go/NoGo task ($n = 2$).

30 31 32 33 34 35 36 37 ***Longitudinal neurophysiological changes associated with mindfulness practice***

38
39
40 In healthy individuals, learning mindfulness was associated with decreased alpha amplitude
41 ($n = 2$ studies), increased ($n = 1$) as well as decreased ($n = 1$) theta amplitude, and changes
42 in asymmetry with an increase relative left-sided activation ($n = 1$).

43
44
45
46
47 In participants with chronic pain, a course of mindfulness was associated with a decrease in
48 beta amplitude ($n = 1$). In patients with depression and suicidal ideation, a relative increase
49 in left-sided activation following mindfulness training was observed, while the inverse pattern
50 with a relative decrease in left-sided activation was reported in patients in remission from
51 depression.

Running title: The neurophysiology of mindfulness

Discussion

1
2
3 The main finding to emerge from the systematic review is an increase in alpha power
4 associated with mindfulness relative to a resting state. Additional effects have been reported
5 in the oscillation bandwidths, including a majority trend towards increased theta power
6 during meditation compared to resting state. The patterns of increased alpha and theta
7 amplitude associated with meditation were observed in both experienced and novice
8 meditators. Clinical studies of mindfulness-based interventions revealed a shift towards
9 greater relative left-sided activation which may be associated with increased positive affect.
10 However, these findings have been mixed with reports of increases, decreases as well as no
11 differences, particularly in other bandwidths, but also in alpha and theta bandwidths.
12
13
14
15
16
17
18
19
20
21
22
23

24 Alpha synchronization has been regarded as one of the 'signatures' of meditation as it has
25 been consistently observed across a range of different meditation practices relatively
26 independent of both technique and degree of practice (Fell et al., 2010). In the present
27 review, increased alpha synchronization during meditation as compared to a resting state
28 was reported 65% of papers that analyzed this outcome (12 out of 18), all of which involved
29 healthy participants, including both novice (Lo et al., 2003; Milz et al., 2014; Takahashiet al.,
30 2005; Yu et al., 2011) and experienced meditators (Ahani et al., 2014; Arita, 2012; Cahn et
31 al., 2013; Dunn et al., 1999; Hinterberger et al., 2014; Huang and Lo, 2009; Kasamatsu and
32 Hirai, 1966; Lagopoulos et al., 2009). Most of the studies had examined participants during
33 mindfulness in comparison to a resting state with eyes closed with a few exceptions (ex.
34 Takahashiet al., 2005). However, the findings have not been wholly consistent as a few
35 studies found no differences with mindfulness in novice (Kubota et al., 2001) or experienced
36 (Cahn et al., 2010; Lehmann et al., 2012) participants, as well as decreased alpha power
37 during mindfulness (Amihai and Kozhevnikov, 2014). It is of note that none of the studies
38 involving clinical populations had analyzed or reported findings on alpha power.
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56

57 Comparisons of mindfulness with performance on attention tasks reported no differences in
58 alpha power with eyes closed while attending to auditory clicks (Becker and Shapiro, 1981);
59
60
61
62
63
64
65

Running title: The neurophysiology of mindfulness

1
2 with a time production task (Berkovich-Ohana et al., 2013); and with an eyes-open session
3 watching a video about neurofeedback (Stinson and Arthur, 2013), although Ren et al.
4 (2011) found lower levels of alpha compared to a problem-solving task.
5
6

7 The functional significance of alpha has been much debated. Alpha synchronization has
8 been understood as reflecting the 'de-activation' of cortical areas as a signifier of the brain
9 'idling' since it occurs during relaxed eyes closed wakefulness (Shaw, 1996; Pfurtscheller et
10 al., 1996). The increase in alpha synchronization with mindfulness as compared to an eyes
11 closed rest may indicate even greater levels of synchronization associated with mindfulness.
12 According to the 'brain idling' hypothesis, the effect suggests that meditation generates
13 greater cortical de-activation than during an eyes closed resting state. However, Shaw
14 (1996) proposes that there is a paradoxical response which distinguishes between 'outer-
15 directed' and 'inner-directed' attention. While 'outer-directed' attention is associated with
16 alpha desynchronisation, 'inner-directed' attention, which is also referred to as 'intention,' is
17 associated with increases in alpha power. In support, tasks requiring memory (Jensen et al.,
18 2002) and imagination (Cooper et al., 2006) lead to increases in alpha power. Mindfulness
19 improves the training and development of various attention networks (sustained, executive,
20 executive, selective, and re-orienting) in terms of its focused-attention aspects and
21 awareness in terms of its open-monitoring aspects (Lutz et al., 2008; Vago and Silbersweig,
22 2012). As such, it is possible to infer that increased alpha power associated with
23 mindfulness is evidence that alpha synchronization is indeed a signifier of increased
24 processing in these various attention modalities (e.g., as per Vago and Silbersweig's (2012)
25 model) with respect to internally generated stimuli.
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50

51 With regards to beta oscillations, of the 12 studies which compared beta activity in
52 meditation with eyes closed rest in healthy individuals, only 3 studies reported that beta
53 amplitude was higher in meditation, involving experienced meditators (Ahani et al., 2014;
54 Cahn et al., 2013) and novices (Dunn et al., 1999). Five studies found no significant
55 differences in experienced practitioners (Cahn et al., 2010; Lagopoulos et al., 2009;
56
57
58
59
60
61
62
63
64
65

Running title: The neurophysiology of mindfulness

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Lehmann et al., 2012) and in novices (Milz et al., 2014; Yu et al., 2011), while one study observed lower beta amplitude in meditation in experienced practitioners (Amihai and Kozhevnikov, 2014), and 5 studies found no significant differences (Cahn et al., 2010; Lagopoulos et al., 2009; Lehmann et al., 2012; Milz et al., 2014; Yu et al., 2011). A comparison of mindfulness with task performance, an eyes open session watching a video about neurofeedback, reported lower amplitude in meditation relative to the task (Stinson and Arthur, 2013). Only one paper reported on beta power in clinical populations, observing pre-post longitudinal decreases in beta power during the resting state which was linked to the practice of mindfulness (Howells et al., 2012).

Interpretations of the significance of beta are mixed because it has been proposed to reflect a reduction in cortical activity as it is associated with barbiturates and benzodiazepines use (Herning et al., 1994), but beta activity has also been attenuated with increasing cognitive task demands (Ray and Cole, 1985) while around 20% of patients with deficit hyperactivity disorder exhibit 'excessive' beta activity, which is associated with elevated behavioural problems (Clarke et al., 2001).

Increased theta power has been considered to be another key feature of meditation (Josipovic, 2010; Fell et al., 2010). This pattern was to some extent borne out in the present review and was observed in both novice and experienced meditators, although there did appear to be a slight weighting towards this effect being more prevalent in experienced practitioners. Of the 19 studies that compared theta activity in meditation with eyes closed rest, a majority ($n = 11$) reported that theta power was higher in mindfulness, including 8 with experienced practitioners (Ahani et al., 2014; Arita, 2012; Cahn et al., 2010; Chan et al., 2008; Kasamatsu and Hirai, 1966; Lagopoulos et al., 2009; Lomas et al., 2014), but only 2 with novices (Kubota et al., 2001; Takahashiet al., 2005), plus also Tanaka et al. (2014), who found this effect with both novice and experienced practitioners. Against this, 3 studies reported that theta was lower during mindfulness compared to eyes-closed rest, 2 of which involved novices (Dunn et al., 1999; Yu et al., 2011) and 1 involving experienced

Running title: The neurophysiology of mindfulness

1
2 practitioners (Huang and Lo, 2009). Moreover, 2 studies found no significant differences in
3 experienced (Amihai and Kozhevnikov, 2014) and novice practitioners (Milz et al., 2014). An
4 additional 2 longitudinal studies also observed pre-post decreases in theta power during the
5 resting state which was linked to the practice of mindfulness (Saggar et al., 2012; Tang et
6 al., 2009). Only one paper reported on theta power in clinical populations, observing pre-
7 post longitudinal increases in theta power (during the resting state) linked to the practice of
8 mindfulness (Howells et al., 2012).
9

10
11 The presence of theta along with alpha synchronization during mindfulness lends support to
12 the hypothesis that increased alpha power during signifies internalized attention rather than
13 the brain 'idling' because theta synchronization is widely viewed as a marker of executive
14 functioning. Theta activity has been linked to various types of cognitive activity, including
15 switching and orienting attention (Dietl et al., 1999), processing of new information
16 (Grunwald et al., 1999), and memory in episodic encoding and retrieval (Klimesch et al.,
17 1997), and theta power increases as task demands increase (Klimesch et al., 1997). Taken
18 together, the finding suggests that mindfulness constitutes a state of enhanced internally-
19 directed attention. Theta oscillations during wakefulness generally occur maximally in the
20 frontal-midline regions of the brain, particularly in the prefrontal cortex (Asada et al., 1999)
21 and may be localized to the anterior cingulate cortex (Onton et al., 2005), in contrast to theta
22 activity during REM sleep, which is generated mainly by the hippocampus (Cantero et al.,
23 2003). These regions are centrally involved in the executive control of attention, as well as
24 other higher-level cognitive activities such as volition and planning (Posner and Dehaene,
25 1994; Miller and Cohen, 2001), and have been proposed as central to the development of
26 attention and awareness in meditation (Newberg and Iversen, 2003).
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51

52
53 This interpretation is strengthened by the differences observed between experienced and
54 novice practitioners, in which the former were more reliably found across the studies to
55 exhibit higher levels of theta activation during meditation in comparison to a resting state,
56 suggesting that enhanced theta activation during meditation is to some extent a function of
57
58
59
60
61
62
63
64
65

Running title: The neurophysiology of mindfulness

training and practice in meditation by learning to maintain an inner-directed attention.

Furthermore, it has been suggested that the co-presence of theta and alpha in mindfulness indicates a state of 'relaxed alertness' (Britton et al., 2014), which is corroborated by qualitative self-reports of their experiences in mindfulness (Cahn and Polich, 2006).

Fewer studies have reported delta and gamma activity with mixed findings and have all been limited to healthy individuals. Slow wave delta band activity is more commonly associated with sleep, particularly during deep non-REM stages (Hofle et al., 1997). It has been suggested though that an increase in delta activity during wakefulness reflects attention to internal processing during the performance of cognitive tasks, such as difficult arithmetical calculation tasks (Harmony et al., 1996). The reports of delta activity have generally found no differences (Lagopoulos et al., 2009; Amihai and Kozhevnikov, 2014; Milz et al., 2014), although reduced (Dunn et al., 1999) as well as increased amplitudes, which were localized to frontal regions (Cahn et al., 2010), have been described in comparison to an eyes closed resting state in novice (Dunn et al., 1999; Milz et al., 2014) and experienced (Lagopoulos et al., 2009; Cahn et al., 2010; Amihai and Kozhevnikov, 2014) meditators. Stinson and Arthur (2013) also found lower amplitude during meditation compared a control task of watching a neurofeedback video.

Gamma synchronization is purported to reflect activity in the default mode network (Berkovich-Ohana et al., 2012) which refers to the self-referential and reflective thoughts that occur in the absence of requirements to respond to external stimuli (Buckner, Andrews-Hanna, and Schacter, 2008). With mindfulness, gamma power has been reported as increased (Berkovich-Ohana et al., 2012; Cahn et al., 2010; Lehmann et al., 2012) as well as showing no differences (Amihai and Kozhevnikov, 2014; Milz et al., 2014) in comparison with an eyes closed resting state. Of interest, increased gamma activity was observed in experienced meditators (Berkovich-Ohana et al., 2012; Cahn et al., 2010; Lehmann et al., 2012), although no differences were also found in both experienced (Amihai and Kozhevnikov, 2014) and novice (Milz et al., 2014) meditators. In comparison with a control

Running title: The neurophysiology of mindfulness

1 task, lower amplitude was reported during mindfulness as compared to a neurofeedback
2 video task (Stinson and Arthur, 2013). In addition, studying experienced Zen meditators,
3 Hauswald et al. (2015) found that gamma power during meditation correlated both with
4 levels of trait mindfulness and years of meditation practice. Ferrarelli et al. (2013) also
5 reported a correlation between meditation experience and gamma power during non-REM
6 sleep, but Berkovich-Ohana et al. (2012) found no difference in coherence between
7 meditation and rest. Gamma oscillations have also been implicated in theories of
8 consciousness, in which the fast rhythmic synchronization of neural discharges provide the
9 necessary spatial and temporal links to bind processing across different brain areas, thereby
10 integrating disparate experiential qualia into a coherent state of moment-to-moment
11 awareness (Singer, 1993; Tallon-Baudry and Bertrand, 1999). Increased gamma power
12 during mindfulness thus might indicate a more unified and coherent mental state.
13
14
15
16
17
18
19
20
21
22
23
24
25
26

27 In addition to analysis of specific bandwidths, patterns of asymmetric brain activation have
28 been examined in which left prefrontal activity has been associated with positive affect and
29 'approach-related' behavior, and right prefrontal activity with negative affect and 'withdrawal-
30 related' behavior (Davidson, 1992). If mindfulness is associated with enhanced subjective
31 wellbeing, then its practice should be linked to greater left prefrontal activity. Such an
32 asymmetry has been observed during mindfulness in experienced meditators relative to an
33 eyes closed resting state (Amihai and Kozhevnikov, 2014; Chan et al., 2008). Following
34 mindfulness training, similar changes have been reported in novice participants who were
35 healthy volunteers (Davidson et al., 2003) as well as with a history of suicidal ideation
36 (Barnhofer et al., 2007). In novice participants with a history of depression, there have been
37 reports of no differences (Milz et al., 2014), increased (Barnhofer et al., 2010) and
38 decreased (Keune et al., 2011) left-sided activation.
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54

55 Using event-related potentials, reduced P3b in response to distractor stimuli (Slagter et al.,
56 2007) and faster attentional disengagement from a dominant global presentation in order to
57 focus in on specific stimuli (van Leeuwen et al., 2012) was observed in experienced
58
59
60
61
62
63
64
65

Running title: The neurophysiology of mindfulness

1
2 meditators. Likewise, Delgado et al. (2013) found that experienced Vipassana meditators
3 demonstrated larger P3b amplitudes to a *target* tone after meditation than before meditation,
4 findings, which are interpreted as reflecting increased attentional engagement following
5 meditation, given that P3b is interpreted as reflecting allocation of attentional resources to
6 incoming stimulation to facilitate information processing, thus corroborating the notion of
7 mindfulness as a system of attention training. Moreover, anticipatory and pain-evoked ERPs
8 to acute pain were reduced in participants who received mindfulness training but not in
9 controls (Brown and Jones, 2013). Sobolewski et al. (2011) explored the impact of
10 meditation practice on late positive potential (LPP), the amplitude of which tends to be
11 greater in ERPs evoked by emotionally arousing images, particularly ones that are
12 negatively valenced. While control participants with no meditation experience showed an
13 increase in LPP amplitude in response to negative stimuli, no such increases were observed
14 in meditators, suggesting that the latter were less affected by negative emotional load than
15 control participants; in contrast, both groups responded equally to positively-valenced
16 stimuli. Teper and Inzlicht (2014) explored participants' neuroaffective reaction to rewarding,
17 aversive and neutral feedback, as gauged by feedback-related negativity (FRN), a brain
18 response that differentiates positive from negative feedback, reporting that trait levels of
19 mindfulness in novice meditators predicted less differentiation of reward from neutral
20 feedback. Lakey et al. (2011) explored the impact of brief mindfulness training on
21 performance of a P300-based brain-compute interface task. Compared to non-meditating
22 control participants, the experimental subjects produced significantly larger P300 amplitudes
23 and were also more accurate at the task which was understood as suggesting that the
24 experimental participants were better able to harness present-moment attentional resources.
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50

51 Working with patients with ADHD, Schoenberg et al. (2014) explored the impact of
52 Mindfulness-Based Cognitive Therapy (MBCT) on error processing (ERN, Pe), conflict
53 monitoring (NoGo-N2), and inhibitory control (NoGo-P3) in relation to a continuous
54 performance task (CPT-X). Compared to matched controls, MBCT was linked to increased
55
56
57
58
59
60
61
62
63
64
65

Running title: The neurophysiology of mindfulness

1
2 Pe and NoGo-P3 amplitudes, which coincided with reduced 'hyperactivity/impulsivity' and
3 'inattention' symptomatology. In a trial involving patients currently diagnosed with major
4 depressive disorder, Schoenberg and Speckens (2014) found that an MBCT intervention
5 had a modulating effect on evoked FM-theta power during a Go/NoGo task: enhanced
6 event-related synchronization (ERS) in the late temporal window was observed pre-to-post
7 for the experimental group, with the reverse pattern found in control participants. It was
8 suggested that these findings were reflective of optimized allocation of attentional resources
9 as a result of the intervention. Moreover, these modulated ERS dynamics were also found to
10 correlate with ameliorated depressive and rumination symptoms in the MBCT group.

11
12 Studying patients with chronic pain at risk of opioid abuse, Garland et al. (2015) found that a
13 Mindfulness-Oriented Recovery Enhancement intervention was able to enhance natural
14 reward processing. In particular, the intervention was associated with increases in LPP in
15 response to natural reward stimuli relative to neutral stimuli, which also correlated with
16 reduced opioid craving from pre- to post-treatment. Jo et al. (2014) explored the Readiness
17 Potential correlates of the intentional binding effect, and found that early neural activity
18 correlates with the participants' reports of initiating a voluntary action; however, there were
19 no differences between experienced Zen meditators and matched controls in this regard.

20
21 A significant limitation of the present systematic review has been the variability of the
22 measures which were acquired and reported such that a meta-analysis was not feasible for
23 any of the measures because there were no more than 3 studies which used the same
24 measure at the same site. The quality was assessed for each of the studies using the
25 Quality Assessment Tool for Quantitative Studies (National Collaborating Centre for Methods
26 and Tools, 2008), revealing considerable variation. Clinical studies were generally of higher
27 quality as they tended to keep track of withdrawal and attrition rates and used standardized
28 meditation protocols. Furthermore, a key issue was limited reporting on participants' prior
29 level of meditation experience. Some studies reported this in terms of years, some in terms
30 of total number of hours, and a few omitted to specify this. Moreover, there was variation in
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Running title: The neurophysiology of mindfulness

1
2 the criteria for which studies rated participants as 'experienced'; in terms of years, this
3 ranged from 1 year (Kasamatsu and Hirai, 1966) to 9 years (Lagopoulos et al., 2009), while
4 in terms of hours this ranged from 40 hours (Hinterberger et al., 2011) to 1740 hours
5 (Berkovich-Ohana et al., 2012). We applied the lowest of these cutoffs such that an
6 'experienced' (i.e., non-novice) meditator was considered to have been meditating for longer
7 than 1 year or have completed more than 40 hours of meditation. Arguably hours would be
8 a better metric than years since it better reflects a person's general amount of practice;
9 however, it is recommended that future studies report both hours and years which would
10 provide some indication of the 'intensity' of participants' practice. Another issue was key
11 poor and/or inconsistent reporting on the nature of participants' meditation practice.
12

13 Although all the studies included in the review featured mindfulness specifically (or a
14 functional equivalent), even this is a somewhat generic label, with nuances and differences
15 among practices that can be classified as such mindfulness prior level of meditation
16 experience. Many studies had not described in detail the form and type of mindfulness
17 practice engaged in by participants.
18

19 In conclusion, the burgeoning literature on EEG investigations of mindfulness is beginning to
20 highlight some consistent trends, most notably with respect to increased amplitude in the
21 alpha and theta bandwidths. The co-presence of elevated alpha and theta waves may
22 reflect a state of 'relaxed alertness as alpha and theta can both be interpreted as signifiers of
23 increased attention with alpha specifically representing internalized attention and both have
24 also been identified as indexing states of relaxation. Further work will be needed to explore
25 the nuances of brain states associated with mindfulness, particularly with respect to the
26 other bandwidths and measures such as ERP and asymmetry, to elucidate the differences
27 between mindfulness and other meditation practices, and to further explore the impact of
28 factors such as degree of meditation practice.
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Running title: The neurophysiology of mindfulness

References

- 1
2
3 Aftanas, L. I., & Golocheikine, S. A. (2001). Human anterior and frontal midline theta and
4 lower alpha reflect emotionally positive state and internalized attention: High-resolution EEG
5 investigation of meditation. *Neuroscience Letters*, *310*(1), 57-60. doi:
6
7 [http://dx.doi.org/10.1016/S0304-3940\(01\)02094-8](http://dx.doi.org/10.1016/S0304-3940(01)02094-8)
8
9
10
11 Ahani, A., Wahbeh, H., Nezamfar, H., Miller, M., Erdogmus, D., & Oken, B. (2014).
12 Quantitative change of EEG and respiration signals during mindfulness meditation. *Journal*
13 *of NeuroEngineering and Rehabilitation*, *11*(87), 10.1186.
14
15
16
17
18 Amihai, I., & Kozhevnikov, M. (2014). Arousal vs. relaxation: A comparison of the
19 neurophysiological and cognitive correlates of Vajrayana and Theravada meditative
20 practices. *PLoS One*. doi: 10.1371/journal.pone.0102990
21
22
23
24
25 Arita, H. (2012). Anterior prefrontal cortex and serotonergic system activation during Zen
26 meditation practice induces negative mood improvement and increased alpha band in EEG.
27 *Rinsho Shinkeigaku (Clinical Neurology)*, *52*(11), 1279-1280. doi:
28
29 [10.5692/clinicalneuro.52.1279](http://dx.doi.org/10.5692/clinicalneuro.52.1279)
30
31
32
33
34 Asada, H., Fukuda, Y., Tsunoda, S., Yamaguchi, M., & Tonoike, M. (1999). Frontal midline
35 theta rhythms reflect alternative activation of prefrontal cortex and anterior cingulate cortex in
36 humans. *Neuroscience Letters*, *274*(1), 29-32. doi: 10.1016/S0304-3940(99)00679-5
37
38
39
40
41 Barnhofer, T., Chittka, T., Nightingale, H., Visser, C., & Crane, C. (2010). State effects of two
42 forms of meditation on prefrontal EEG asymmetry in previously depressed individuals.
43 *Mindfulness*, *1*(1), 21-27. doi: 10.1007/s12671-010-0004-7
44
45
46
47
48 Barnhofer, T., Duggan, D., Crane, C., Hepburn, S., Fennell, M. J., & Williams, J. M. (2007).
49 Effects of meditation on frontal alpha-asymmetry in previously suicidal individuals.
50 *Neuroreport*, *18*(7), 709-712. doi: 10.1097/WNR.0b013e3280d943cd
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Running title: The neurophysiology of mindfulness

1
2 Berkovich-Ohana, A., Glicksohn, J., & Goldstein, A. (2012). Mindfulness-induced changes in
3 gamma band activity – Implications for the default mode network, self-reference and
4 attention. *Clinical Neurophysiology*, 123(4), 700-710. doi:
5

6 <http://dx.doi.org/10.1016/j.clinph.2011.07.048>
7

8
9 Berkovich-Ohana, A., Glicksohn, J., & Goldstein, A. (2013). Studying the default mode and
10 its mindfulness-induced changes using EEG functional connectivity. *Social Cognitive and*
11 *Affective Neuroscience*. doi: 10.1093/scan/nst153
12
13

14
15 Bostanov, V., Keune, P. M., Kotchoubey, B., & Hautzinger, M. (2012). Event-related brain
16 potentials reflect increased concentration ability after mindfulness-based cognitive therapy
17 for depression: a randomized clinical trial. *Psychiatry Research*, 199(3), 174-180. doi:
18
19

20 [10.1016/j.psychres.2012.05.031](http://dx.doi.org/10.1016/j.psychres.2012.05.031)
21
22

23
24 Bowen, S., Witkiewitz, K., Clifasefi, S. L., Grow, J., Chawla, N., Hsu, S. H., . . . Larimer, M.
25 E. (2014). Relative efficacy of mindfulness-based relapse prevention, standard relapse
26 prevention, and treatment as usual for substance use disorders: A randomized clinical trial.
27
28
29
30
31 *JAMA Psychiatry*, 71(5), 547-556. doi: 10.1001/jamapsychiatry.2013.4546
32

33
34 Brown, C. A., & Jones, A. K. (2010). Meditation experience predicts less negative appraisal
35 of pain: Electrophysiological evidence for the involvement of anticipatory neural responses.
36
37
38 *Pain*, 150(3), 428-438. doi: 10.1016/j.pain.2010.04.017
39

40
41 Brown, C. A., & Jones, A. K. (2013). Psychobiological correlates of improved mental health
42 in patients with musculoskeletal pain after a mindfulness-based pain management program.
43
44
45 *Clinical Journal of Pain*, 29(3), 233-244. doi: 10.1097/AJP.0b013e31824c5d9f
46

47
48 Britton, W. B., Lindahl, J. R., Cahn, B. R., Davis, J. H., & Goldman, R. E. (2014). Awakening
49 is not a metaphor: The effects of Buddhist meditation practices on basic wakefulness.
50
51
52 *Annals of the New York Academy of Sciences*, 1307(1), 64-81. doi: 10.1111/nyas.12279
53

54
55 Buckner, R. L., Andrews-Hanna, J. R., & Schacter, D. L. (2008). The brain's default network.
56
57 *Annals of the New York Academy of Sciences*, 1124(1), 1-38. doi: 10.1196/annals.1440.011
58

59
60 Cacioppo, J. T., Tassinary, L. G., & Berntson, G. G. (2007). *Handbook of Psychophysiology*
61 (3rd ed.). Cambridge: Cambridge University Press.
62
63
64
65

Running title: The neurophysiology of mindfulness

1
2 Cahn, B. R., Delorme, A., & Polich, J. (2010). Occipital gamma activation during Vipassana
3 meditation. *Cognitive Processing*, 11(1), 39-56. doi: 10.1007/s10339-009-0352-1

4 Cahn, B. R., Delorme, A., & Polich, J. (2013). Event-related delta, theta, alpha and gamma
5 correlates to auditory oddball processing during Vipassana meditation. *Social, Cognitive and*
6 *Affective Neuroscience*, 8(1), 100-111. doi: 10.1093/scan/nss060

7
8 Cahn, B. R., & Polich, J. (2006). Meditation states and traits: EEG, ERP, and neuroimaging
9 studies. *Psychological Bulletin*, 132(2), 180-211. doi: 10.1037/0033-2909.132.2.180

10
11 Cantero, J. L., Atienza, M., Stickgold, R., Kahana, M. J., Madsen, J. R., & Kocsis, B. (2003).
12 Sleep-dependent {theta} oscillations in the human hippocampus and neocortex. *The Journal*
13 *of Neuroscience*, 23(34), 10897-10903.

14
15 Chalmers, D. J. (1995). Facing up to the problem of consciousness. *Journal of*
16 *Consciousness Studies*, 2, 200-219.

17
18 Chambers, R., Lo, B., & Allen, N. (2008). The impact of intensive mindfulness training on
19 attentional control, cognitive style, and affect. *Cognitive Therapy and Research*, 32(3), 303-
20 322. doi: 10.1007/s10608-007-9119-0

21
22 Chan, A. S., Han, Y. M., & Cheung, M. C. (2008). Electroencephalographic (EEG)
23 measurements of mindfulness-based Triarchic body-pathway relaxation technique: A pilot
24 study. *Applied Psychophysiology and Biofeedback*, 33(1), 39-47. doi: 10.1007/s10484-008-
25 9050-5

26
27 Chiesa, A., Calati, R., & Serretti, A. (2011). Does mindfulness training improve cognitive
28 abilities? A systematic review of neuropsychological findings. *Clinical Psychology Review*,
29 31(3), 449-464. doi: 10.1016/j.cpr.2010.11.003

30
31 Clarke, A. R., Barry, R. J., McCarthy, R., & Selikowitz, M. (2001). Excess beta activity in
32 children with attention-deficit/hyperactivity disorder: An atypical electrophysiological group.
33 *Psychiatry Research*, 103(2), 205-218. doi: 10.1016/S0165-1781(01)00277-3

34
35 Cooper, N. R., Burgess, A. P., Croft, R. J., & Gruzelier, J. H. (2006). Investigating evoked
36 and induced electroencephalogram activity in task-related alpha power increases during an
37 internally directed attention task. *Neuroreport*, 17(2), 205-208.

Running title: The neurophysiology of mindfulness

1
2 Cousins, L. S. (1996). The dating of the historical Buddha: A review article. *Journal of the*
3 *Royal Asiatic Society (Third Series)*, 6(1), 57-63. doi: doi:10.1017/S1356186300014760

4 Davidson, R. J. (1992). Anterior cerebral asymmetry and the nature of emotion. *Brain and*
5 *Cognition*, 20(1), 125-151. doi: 10.1016/0278-2626(92)90065-T

6
7
8 Davidson, R. J., Kabat-Zinn, J., Schumacher, J., Rosenkranz, M., Muller, D., Santorelli, S.
9 F., . . . Sheridan, J. F. (2003). Alterations in brain and immune function produced by

10
11 mindfulness meditation. *Psychosomatic Medicine*, 65(4), 564-570. doi:

12
13 10.1097/01.psy.0000077505.67574.e3

14
15
16 Delgado-Pastor, L. C., Perakakis, P., Subramanya, P., Telles, S., & Vila, J. (2013).

17
18 Mindfulness (Vipassana) meditation: Effects on P3b event-related potential and heart rate
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

10.1016/j.ijpsycho.2013.07.006

51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

10.1023/A:1023498629385
Dunn, B. R., Hartigan, J. A., & Mikulas, W. L. (1999). Concentration and mindfulness
meditations: Unique forms of consciousness? *Applied Psychophysiology and Biofeedback*,
24(3), 147-165. doi: 10.1023/A:1023498629385

Egan, H. D. (1978). Christian apophatic and kataphatic mysticisms. *Theological Studies*,
39(3), 399-426. doi: 10.1177/004056397803900301

Fell, J. (2004). Identifying neural correlates of consciousness: The state space approach.
Consciousness and Cognition, 13(4), 709-729. doi: 10.1016/j.concog.2004.07.001

Fell, J., Axmacher, N., & Haupt, S. (2010). From alpha to gamma: Electrophysiological
correlates of meditation-related states of consciousness. *Medical Hypotheses*, 75(2), 218-
224. doi: <http://dx.doi.org/10.1016/j.mehy.2010.02.025>

Ferrarelli, F., Smith, R., Dentico, D., Riedner, B. A., Zennig, C., Benca, R. M., . . . Tononi, G.
(2013). Experienced mindfulness meditators exhibit higher parietal-occipital EEG gamma
activity during NREM sleep. *PLoS One*, 8(8), e73417. doi: 10.1371/journal.pone.0073417

Running title: The neurophysiology of mindfulness

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Fresco, D. M., Moore, M. T., van Dulmen, M. H. M., Segal, Z. V., Ma, S. H., Teasdale, J. D., & Williams, J. M. G. (2007). Initial psychometric properties of the experiences questionnaire: Validation of a self-report measure of decentering. *Behavior Therapy, 38*(3), 234-246. doi: 10.1016/j.beth.2006.08.003

Garland, E., Froeliger, B., & Howard, M. (2015). Neurophysiological evidence for remediation of reward processing deficits in chronic pain and opioid misuse following treatment with Mindfulness-Oriented Recovery Enhancement: exploratory ERP findings from a pilot RCT. *Journal of Behavioral Medicine, 38*(2), 327-336. doi: 10.1007/s10865-014-9607-0

Garland, E. L., Manusov, E. G., Froeliger, B., Kelly, A., Williams, J. M., & Howard, M. O. (2014). Mindfulness-oriented recovery enhancement for chronic pain and prescription opioid misuse: Results from an early-stage randomized controlled trial. *Journal of Consulting and Clinical Psychology, 82*(3), 448-459. doi: 10.1037/a0035798

Grunwald, M., Weiss, T., Krause, W., Beyer, L., Rost, R., Gutberlet, I., & Gertz, H.-J. (1999). Power of theta waves in the EEG of human subjects increases during recall of haptic information. *Neuroscience Letters, 260*(3), 189-192. doi: [http://dx.doi.org/10.1016/S0304-3940\(98\)00990-2](http://dx.doi.org/10.1016/S0304-3940(98)00990-2)

Harmony, T., Fernández, T., Silva, J., Bernal, J., Díaz-Comas, L., Reyes, A., . . . Rodríguez, M. (1996). EEG delta activity: An indicator of attention to internal processing during performance of mental tasks. *International Journal of Psychophysiology, 24*(1-2), 161-171. doi: [http://dx.doi.org/10.1016/S0167-8760\(96\)00053-0](http://dx.doi.org/10.1016/S0167-8760(96)00053-0)

Hauswald, A., Übelacker, T., Leske, S., & Weisz, N. (2015). What it means to be Zen: Marked modulations of local and interareal synchronization during open monitoring meditation. *Neuroimage, 108*, 265-273. doi: 10.1016/j.neuroimage.2014.12.065

Herning, R. I., Glover, B. J., Koepl, B., Phillips, R. L., & London, E. D. (1994). Cocaine-induced increases in EEG alpha and beta activity: evidence for reduced cortical processing. *Neuropsychopharmacology, 11*(1), 1-9. doi: 10.1038/npp.1994.30

Running title: The neurophysiology of mindfulness

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Hinterberger, T., Kamei, T., & Walach, H. (2011). Psychophysiological classification and staging of mental states during meditative practice. *Biomedizinische Technik (Biomedical Engineering)*, *56*(6), 341-350. doi: 10.1515/bmt.2011.021

Hinterberger, T., Schmidt, S., Kamei, T., & Walach, H. (2014). Decreased electrophysiological activity represents the conscious state of emptiness in meditation. *Frontiers in Psychology*, *5*, 99. doi: 10.3389/fpsyg.2014.00099

Hofle, N., Paus, T., Reutens, D., Fiset, P., Gotman, J., Evans, A. C., & Jones, B. E. (1997). Regional cerebral blood flow changes as a function of delta and spindle activity during slow wave sleep in humans. *The Journal of Neuroscience*, *17*(12), 4800-4808.

Hofmann, S. G., Sawyer, A. T., Witt, A. A., & Oh, D. (2010). The effect of mindfulness-based therapy on anxiety and depression: A meta-analytic review. *Journal of Consulting and Clinical Psychology*, *78*(2), 169-183. doi: 10.1037/a0018555

Howells, F. M., Ives-Deliperi, V. L., Horn, N. R., & Stein, D. J. (2012). Mindfulness based cognitive therapy improves frontal control in bipolar disorder: a pilot EEG study. *BMC Psychiatry*, *12*, 15. doi: 10.1186/1471-244x-12-15

Huang, H. Y., & Lo, P. C. (2009). EEG dynamics of experienced Zen meditation practitioners probed by complexity index and spectral measure. *Journal of Medical Engineering & Technology*, *33*(4), 314-321. doi: 10.1080/03091900802602677

Jensen, O., Gelfand, J., Kounios, J., & Lisman, J. E. (2002). Oscillations in the alpha band (9–12 Hz) increase with memory load during retention in a short-term memory task. *Cerebral Cortex*, *12*(8), 877-882. doi: doi:10.1093/cercor/12.8.87

Jha, A. P., Krompinger, J., & Baime, M. J. (2007). Mindfulness training modifies subsystems of attention. *Cognitive, Affective, & Behavioral Neuroscience*, *7*(2), 109-119. doi: 10.3758/CABN.7.2.109

Jo, H.-G., Wittmann, M., Hinterberger, T., & Schmidt, S. (2014). The readiness potential reflects intentional binding. *Frontiers in Human Neuroscience*, *8*, 421. doi: 10.3389/fnhum.2014.00421

Running title: The neurophysiology of mindfulness

1
2 Josipovic, Z. (2010). Duality and nonduality in meditation research. *Consciousness and*
3 *Cognition*, 19(4), 1119-1121. doi: <http://dx.doi.org/10.1016/j.concog.2010.03.016>

4 Kabat-Zinn, J. (1982). An outpatient program in behavioral medicine for chronic pain patients
5 based on the practice of mindfulness meditation: Theoretical considerations and preliminary
6 results. *General Hospital Psychiatry*, 4(1), 33-47. doi: [http://dx.doi.org/10.1016/0163-](http://dx.doi.org/10.1016/0163-8343(82)90026-3)
7
8
9 8343(82)90026-3

10 Kabat-Zinn, J. (2003). Mindfulness-based interventions in context: Past, present, and future.
11
12 *Clinical Psychology: Science and Practice*, 10(2), 144-156. doi: 10.1093/clipsy.bpg016

13 Kasamatsu, A., & Hirai, T. (1966). Electroencephalogram study on the Zen meditation
14 (Zazen). *Psychiatry and Clinical Neurosciences*, 20(4), 315-336. doi: 10.1111/j.1440-
15
16
17 1819.1966.tb02646.x

18 Kerr, C. E., Jones, S. R., Wan, Q., Pritchett, D. L., Wasserman, R. H., Wexler, A., . . . Moore,
19
20
21 C. I. (2011). Effects of mindfulness meditation training on anticipatory alpha modulation in
22 primary somatosensory cortex. *Brain Research Bulletin*, 85(3-4), 96-103. doi:
23
24
25 10.1016/j.brainresbull.2011.03.026

26 Keune, P. M., Bostanov, V., Hautzinger, M., & Kotchoubey, B. (2011). Mindfulness-based
27
28
29 cognitive therapy (MBCT), cognitive style, and the temporal dynamics of frontal EEG alpha
30
31
32 asymmetry in recurrently depressed patients. *Biological Psychology*, 88(2-3), 243-252. doi:
33
34
35 10.1016/j.biopsycho.2011.08.008

36 Keune, P. M., Bostanov, V., Hautzinger, M., & Kotchoubey, B. (2013). Approaching
37
38
39 dysphoric mood: State-effects of mindfulness meditation on frontal brain asymmetry.
40
41
42 *Biological Psychology*, 93(1), 105-113. doi: 10.1016/j.biopsycho.2013.01.016

43 Klimesch, W., Doppelmayr, M., Schimke, H., & Ripper, B. (1997). Theta synchronization and
44
45
46 alpha desynchronization in a memory task. *Psychophysiology*, 34(2), 169-176. doi:
47
48
49 10.1111/j.1469-8986.1997.tb02128.x

50 Kubota, Y., Sato, W., Toichi, M., Murai, T., Okada, T., Hayashi, A., & Sengoku, A. (2001).
51
52
53 Frontal midline theta rhythm is correlated with cardiac autonomic activities during the
54
55
56

Running title: The neurophysiology of mindfulness

1 performance of an attention demanding meditation procedure. *Cognitive Brain Research*,
2 11(2), 281-287. doi: 10.1016/S0926-6410(00)00086-0

3
4 Lagopoulos, J., Xu, J., Rasmussen, I., Vik, A., Malhi, G. S., Eliassen, C. F., . . . Ellingsen, O.
5
6 (2009). Increased theta and alpha EEG activity during nondirective meditation. *Journal of*
7
8 *Alternative and Complementary Medicine*, 15(11), 1187-1192. doi: 10.1089/acm.2009.0113

9
10 Lakey, C. E., Berry, D. R., & Sellers, E. W. (2011). Manipulating attention via mindfulness
11
12 induction improves P300-based brain-computer interface performance. *Journal of Neural*
13
14 *Engineering*, 8(2), 025019. doi: 10.1088/1741-2560/8/2/025019

15
16 Ledesma, D., & Kumano, H. (2009). Mindfulness-based stress reduction and cancer: A
17
18 meta-analysis. *Psycho-Oncology*, 18(6), 571-579. doi: doi:10.1002/pon.1400

19
20
21
22 Lehmann, D., Faber, P. L., Achermann, P., Jeanmonod, D., Gianotti, L. R., & Pizzagalli, D.
23
24 (2001). Brain sources of EEG gamma frequency during volitionally meditation-induced,
25
26 altered states of consciousness, and experience of the self. *Psychiatry Research*, 108(2),
27
28 111-121. doi: 10.1016/S0925-4927(01)00116-0

29
30
31
32 Lehmann, D., Faber, P. L., Tei, S., Pascual-Marqui, R. D., Milz, P., & Kochi, K. (2012).
33
34 Reduced functional connectivity between cortical sources in five meditation traditions
35
36 detected with lagged coherence using EEG tomography. *Neuroimage*, 60(2), 1574-1586.
37
38 doi: 10.1016/j.neuroimage.2012.01.042

39
40
41
42 Light, G. A., Williams, L. E., Minow, F., Sprock, J., Rissling, A., Sharp, R., . . . Braff, D. L.
43
44 (2010). Electroencephalography (EEG) and Event-Related Potentials (ERP's) with human
45
46 participants. *Current Protocols in Neuroscience*, CHAPTER, Unit-6.2524. doi:
47
48 10.1002/0471142301.ns0625s52

49
50
51
52 Lo, P. C., & Chang, C. H. (2013). Spatially Nonlinear Interdependence of Alpha-Oscillatory
53
54 Neural Networks under Chan Meditation. *Evidence Based Complementary and Alternative*
55
56 *Medicine*, 2013, 360371. doi: 10.1155/2013/360371

57
58
59
60 Lo, P. C., Huang, M. L., & Chang, K. M. (2003). EEG alpha blocking correlated with
61
62 perception of inner light during zen meditation. *American Journal of Chinese Medicine*,
63
64 31(4), 629-642. doi: 10.1142/s0192415x03001272

Running title: The neurophysiology of mindfulness

1
2 Lomas, T., Edginton, T., Cartwright, T., & Ridge, D. (2014). Men developing emotional
3 intelligence through meditation? Combining narrative, cognitive, and electroencephalography
4 (EEG) evidence. *Psychology of Men and Masculinity*, 15(2), 213-224. doi:

5
6 10.1037/a0032191

7
8 Lutz, A., Slagter, H. A., Dunne, J. D., & Davidson, R. J. (2008). Attention regulation and
9 monitoring in meditation. *Trends in Cognitive Sciences*, 12(4), 163-169. doi:

10
11 10.1016/j.tics.2008.01.005

12
13 Manna, A., Raffone, A., Perrucci, M. G., Nardo, D., Ferretti, A., Tartaro, A., . . . Romani, G.
14
15 L. (2010). Neural correlates of focused attention and cognitive monitoring in meditation.

16
17
18 *Brain Research Bulletin*, 82(1-2), 46-56. doi: 10.1016/j.brainresbull.2010.03.001

19
20 Miller, E. K., & Cohen, J. D. (2001). An integrative theory of prefrontal cortex function.

21
22
23 *Annual Review of Neuroscience*, 24, 167-202. doi: 10.1146/annurev.neuro.24.1.167

24
25 Milz, P., Faber, P. L., Lehmann, D., Kochi, K., & Pascual-Marqui, R. D. (2014). sLORETA
26
27 intracortical lagged coherence during breath counting in meditation-naïve participants.

28
29
30 *Frontiers in Human Neuroscience*, 8(303). doi: 10.3389/fnhum.2014.00303

31
32 Mirsky, A., Anthony, B., Duncan, C., Ahearn, M., & Kellam, S. (1991). Analysis of the
33
34 elements of attention: A neuropsychological approach. *Neuropsychology Review*, 2(2), 109-
35
36 145. doi: 10.1007/BF01109051

37
38
39 Moher, D., Liberati, A., Tetzlaff, J., & Altman, D. G. (2009). Preferred Reporting Items for
40
41 Systematic Reviews and Meta-Analyses: The PRISMA Statement. *PLoS Medicine*, 6(7),

42
43
44 e1000097. doi: doi:10.1371/journal.pmed.1000097

45
46
47 Moore, A., Gruber, T., Derose, J., & Malinowski, P. (2012). Regular, brief mindfulness
48
49 meditation practice improves electrophysiological markers of attentional control. *Frontiers in*

50
51
52 *Human Neuroscience*, 6, 18. doi: 10.3389/fnhum.2012.00018

53
54 Murata, T., Takahashi, T., Hamada, T., Omori, M., Kosaka, H., Yoshida, H., & Wada, Y.
55
56 (2004). Individual trait anxiety levels characterizing the properties of zen meditation.

57
58
59 *Neuropsychobiology*, 50(2), 189-194. doi: 10.1159/000079113

Running title: The neurophysiology of mindfulness

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

National Collaborating Centre for Methods and Tools (2008). *Quality Assessment Tool for Quantitative Studies (QATQS)*. Hamilton, ON: McMaster University.

Newberg, A. B., & Iversen, J. (2003). The neural basis of the complex mental task of meditation: neurotransmitter and neurochemical considerations. *Medical Hypotheses*, *61*(2), 282-291. doi: [http://dx.doi.org/10.1016/S0306-9877\(03\)00175-0](http://dx.doi.org/10.1016/S0306-9877(03)00175-0)

Onton, J., Delorme, A., & Makeig, S. (2005). Frontal midline EEG dynamics during working memory. *Neuroimage*, *27*(2), 341-356. doi: 10.1016/j.neuroimage.2005.04.014

Pasquini, H. A., Tanaka, G. K., Basile, L. F. H., Velasques, B., Lozano, M. D., & Ribeiro, P. (2015). Electrophysiological correlates of long-term Soto Zen meditation. *BioMed Research International*. doi: 10.1155/2015/598496

Pfurtscheller, G., Stancák, A., & Neuper, C. (1996). Event-related synchronization (ERS) in the alpha band — an electrophysiological correlate of cortical idling: A review. *International Journal of Psychophysiology*, *24*(1-2), 39-46. doi: 10.1016/S0167-8760(96)00066-9

Posner, M. I., & Dehaene, S. (1994). Attentional networks. *Trends in Neurosciences*, *17*(2), 75-79. doi: 10.1016/0166-2236(94)90078-7

Posner, M. I., & Petersen, S. E. (1990). The attention system of the human brain. *Annual Review of Neuroscience*, *13*(1), 25-42. doi: doi:10.1146/annurev.ne.13.030190.000325

Raffone, A., & Srinivasan, N. (2010). The exploration of meditation in the neuroscience of attention and consciousness. *Cognitive Processing*, *11*(1), 1-7. doi: 10.1007/s10339-009-0354-z

Rangaswamy, M., Porjesz, B., Chorlian, D. B., Wang, K., Jones, K. A., Kuperman, S., . . .

Reich, T. (2004). Resting EEG in offspring of male alcoholics: beta frequencies. *International Journal of Psychophysiology*, *51*(3), 239-251. doi: 10.1016/j.ijpsycho.2003.09.003

Ray, W. J., & Cole, H. W. (1985). EEG alpha activity reflects attentional demands, and beta activity reflects emotional and cognitive processes. *Science*, *228*(4700), 750-752. doi: 10.1126/science.3992243

Running title: The neurophysiology of mindfulness

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- Ren, J., Huang, Z., Luo, J., Wei, G., Ying, X., Ding, Z., . . . Luo, F. (2011). Meditation promotes insightful problem-solving by keeping people in a mindful and alert conscious state. *Sci China Life Sci*, *54*(10), 961-965. doi: 10.1007/s11427-011-4233-3
- Sadaghiani, S., Scheeringa, R., Lehongre, K., Morillon, B., Giraud, A.-L., D'Esposito, M., & Kleinschmidt, A. (2012). Alpha-band phase synchrony is related to activity in the fronto-parietal adaptive control network. *The Journal of Neuroscience*, *32*(41), 14305-14310.
- Sagar, M., King, B. G., Zanesco, A. P., Maclean, K. A., Aichele, S. R., Jacobs, T. L., . . . Saron, C. D. (2012). Intensive training induces longitudinal changes in meditation state-related EEG oscillatory activity. *Front Hum Neurosci*, *6*, 256. doi: 10.3389/fnhum.2012.00256
- Schmidt, S., Simshäuser, K., Aickin, M., Lüking, M., Schultz, C., & Kaube, H. (2010). Mindfulness-based stress reduction is an effective intervention for patients suffering from migraine—Results from a controlled trial. *European Journal of Integrative Medicine*, *2*(4), 196-196. doi: 10.1016/j.eujim.2010.09.052
- Schoenberg, P. L. A., Hepark, S., Kan, C. C., Barendregt, H. P., Buitelaar, J. K., & Speckens, A. E. M. (2014). Effects of mindfulness-based cognitive therapy on neurophysiological correlates of performance monitoring in adult attention-deficit/hyperactivity disorder. *Clinical Neurophysiology*, *125*(7), 1407-1416. doi: <http://dx.doi.org/10.1016/j.clinph.2013.11.031>
- Schoenberg, P. L., & Speckens, A. E. (2014). Modulation of induced frontocentral theta (Fm- θ) event-related (de-) synchronisation dynamics following mindfulness-based cognitive therapy in Major Depressive Disorder. *Cognitive Neurodynamics*, *8*(5), 373-388. doi: 10.1007/s11571-014-9294-0
- Schoenberg, P. L., & Speckens, A. E. (2015). Multi-dimensional modulations of α and γ cortical dynamics following mindfulness-based cognitive therapy in Major Depressive Disorder. *Cognitive Neurodynamics*, *9*(1), 13-29. doi: 10.1007/s11571-014-9308-y
- Segal, Z. V., Williams, J. M. G., & Teasdale, J. D. (2002). Mindfulness-Based Cognitive Therapy for Depression: A New Approach to Preventing Relapse. New York: Guilford Press.

Running title: The neurophysiology of mindfulness

1 Shapiro, S. L., Astin, J. A., Bishop, S. R., & Cordova, M. (2005). Mindfulness-based stress
2 reduction for health care professionals: Results from a randomized trial. *International Journal*
3 *of Stress Management*, 12(2), 164-176. doi: 10.1037/1072-5245.12.2.164

4
5
6 Shaw, J. C. (1996). Intention as a component of the alpha-rhythm response to mental
7 activity. *International Journal of Psychophysiology*, 24(1-2), 7-23. doi:
8
9 [http://dx.doi.org/10.1016/S0167-8760\(96\)00052-9](http://dx.doi.org/10.1016/S0167-8760(96)00052-9)

10
11
12 Singer, W. (1993). Synchronization of cortical activity and its putative role in information
13 processing and learning. *Annual Review of Physiology*, 55(1), 349-374. doi:
14
15
16
17
18 10.1146/annurev.ph.55.030193.002025

19
20 Slagter, H. A., Lutz, A., Greischar, L. L., Francis, A. D., Nieuwenhuis, S., Davis, J. M., &
21
22 Davidson, R. J. (2007). Mental training affects distribution of limited brain resources. *PLoS*
23 *Biology*, 5(6), e138. doi: 10.1371/journal.pbio.0050138

24
25
26 Slagter, H. A., Lutz, A., Greischar, L. L., Nieuwenhuis, S., & Davidson, R. J. (2009). Theta
27 phase synchrony and conscious target perception: impact of intensive mental training.
28
29
30
31 *Journal of Cognitive Neuroscience*, 21(8), 1536-1549. doi: 10.1162/jocn.2009.21125

32
33 Sobolewski, A., Holt, E., Kublik, E., & Wrobel, A. (2011). Impact of meditation on emotional
34 processing - A visual ERP study. *Neuroscience Research*, 71(1), 44-48. doi:
35
36
37
38 10.1016/j.neures.2011.06.002

39
40 Stinson, B., & Arthur, D. (2013). A novel EEG for alpha brain state training,
41
42 neurobiofeedback and behavior change. *Complementary Therapies in Clinical Practice*,
43
44
45 19(3), 114-118. doi: 10.1016/j.ctcp.2013.03.003

46
47 Takahashi, T., Murata, T., Hamada, T., Omori, M., Kosaka, H., Kikuchi, M., . . . Wada, Y.
48
49
50
51 (2005). Changes in EEG and autonomic nervous activity during meditation and their
52 association with personality traits. *International Journal of Psychophysiology*, 55(2), 199-207.
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
doi: 10.1016/j.ijpsycho.2004.07.004

Tallon-Baudry, C., & Bertrand, O. (1999). Oscillatory gamma activity in humans and its role
in object representation. *Trends in Cognitive Sciences*, 3(4), 151-162. doi: 10.1016/S1364-
6613(99)01299-1

Running title: The neurophysiology of mindfulness

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- Tanaka, G. K., Peressutti, C., Teixeira, S., Cagy, M., Piedade, R., Nardi, A. E., . . .
.Velasques, B. (2014). Lower trait frontal theta activity in mindfulness meditators. *Arquivos de Neuro-Psiquiatria*, *72*(9), 687-693. doi: <http://dx.doi.org/10.1590/0004-282X20140133>
- Tang, Y. Y., Ma, Y., Fan, Y., Feng, H., Wang, J., Feng, S., . . . Fan, M. (2009). Central and autonomic nervous system interaction is altered by short-term meditation. *Proceedings of the National Academy of Sciences of the United States of America*, *106*(22), 8865-8870. doi: [10.1073/pnas.0904031106](https://doi.org/10.1073/pnas.0904031106)
- Teper, R., & Inzlicht, M. (2013). Meditation, mindfulness and executive control: The importance of emotional acceptance and brain-based performance monitoring. *Social Cognitive and Affective Neuroscience*, *8*(1), 85-92. doi: [10.1093/scan/nss045](https://doi.org/10.1093/scan/nss045)
- Teper, R., & Inzlicht, M. (2014). Mindful acceptance dampens neuroaffective reactions to external and rewarding performance feedback. *Emotion*, *14*(1), 105-114. doi: [10.1037/a0034296](https://doi.org/10.1037/a0034296)
- Vago, D. R., & Silbersweig, D. A. (2012). Self-awareness, self-regulation, and self-transcendence (S-ART): a framework for understanding the neurobiological mechanisms of mindfulness. *Frontiers in Human Neuroscience*, *6*, 296. doi: [10.3389/fnhum.2012.00296](https://doi.org/10.3389/fnhum.2012.00296)
- van Leeuwen, S., Singer, W., & Melloni, L. (2012). Meditation increases the depth of information processing and improves the allocation of attention in space. *Front Hum Neurosci*, *6*, 133. doi: [10.3389/fnhum.2012.00133](https://doi.org/10.3389/fnhum.2012.00133)
- Walsh, R., & Shapiro, S. L. (2006). The meeting of meditative disciplines and western psychology: A mutually enriching dialogue. *American Psychologist*, *61*(3), 227-239. doi: [10.1037/0003-066X.61.3.227](https://doi.org/10.1037/0003-066X.61.3.227)
- Williams, J. D., & Gruzelier, J. H. (2001). Differentiation of hypnosis and relaxation by analysis of narrow band theta and alpha frequencies. *International Journal of Clinical and Experimental Hypnosis*, *49*(3), 185-206. doi: [10.1080/00207140108410070](https://doi.org/10.1080/00207140108410070)
- Xue, S.-W., Tang, Y.-Y., Tang, R., & Posner, M. I. (2014). Short-term meditation induces changes in brain resting EEG theta networks. *Brain and Cognition*, *87*, 1-6. doi: [10.1016/j.bandc.2014.02.008](https://doi.org/10.1016/j.bandc.2014.02.008)

Running title: The neurophysiology of mindfulness

1 Yu, X., Fumoto, M., Nakatani, Y., Sekiyama, T., Kikuchi, H., Seki, Y., . . . Arita, H. (2011).

2 Activation of the anterior prefrontal cortex and serotonergic system is associated with
3 improvements in mood and EEG changes induced by Zen meditation practice in novices.
4

5
6 *International Journal of Psychophysiology*, 80(2), 103-111. doi:

7
8 10.1016/j.ijpsycho.2011.02.004
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Figure Legends


Figure 1.The Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) Flow

Diagram

Accepted Manuscript

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Figure 1.



Running title: The neurophysiology of mindfulness

Table 1. Demographics of healthy participants

First author	Year	Meditators	Meditators (male)	Controls	Mean age meditators	Mean years meditating	Meditation type	Study type
Davidson	2003	25 (from 32)	6	16	36	0	MBCT	Pre-post
Dunn	1999	9 (from 10)	NR	-	NR	0	FA& MM	Pre-post
Kerr	2011	12 (from 16)	1	6	31	0	MBCT	Pre-post
Lomas	2014	30	30	-	42.3	10.1	Various (inc. MM)	Pre-post
Moore	2012	12 (from 19)	NR	16 (from 23)	36.9	0	MM	Pre-post
Saggar	2012	22 (from 30)	12	22 (from 30)	49.5	Experienced (yrs NR)	MM (retreat)	Pre-post
Slagter	2007	17	17	23	NR	Experienced (yrs NR)	Vipassana (retreat)	Pre-post
Slagter	2009	17	17	23	NR	Experienced (yrs NR)	Vipassana (retreat)	Pre-post
Tang	2009	40	NR	40	NR	0	Mind-body training	Pre-post
Xue	2014	45	29	24	22.9	0	Mind-body training	Pre-post
Ahani	2014	34	6	-	61	0 (6 weeks training)	MM	Non pre-post
Amihai	2014	19	16	-	44.4	7.7	Vipassana	Non pre-post
Arita	2012	15	NA	-	NA	NA	Zen	Non pre-post
Becker	1981	30 (10 Zen)	17 (Zen=8)	10	32.7 (Zen=37.8)	6.5 (Zen=7.5)	Zen, TM & Yoga	Non pre-post
Berkovich-Ohana	2012	36	NR	12	41.7	3,673 (hrs)	MM	Non pre-post
Berkovich-Ohana	2013	36	NR	12	41.7	3,673 (hrs)	MM	Non pre-post
Brown	2010	12	6	15	34	NR	Various (inc. MM)	Non pre-post
Cahn	2010	16	11	-	45.5	20	Vipassana	Non pre-post
Cahn	2013	16	11	-	45.5	20	Vipassana	Non pre-post
Chan	2008	19	8	-	19-22 (range)	NR	Triarchic	Non pre-post
Delgado	2013	10	10	0	20-61 (range)	7.5	Vipassana	Non pre-post
Ferrarelli	2013	29	14	29	50.7	15.6	MM	Non pre-post
Hauswald	2015	11	5	-	50	12	Zen	Non pre-post
Hinterberger	2011	49	33	-	45	40-1000 (hrs; range)	Various (inc. MM)	Non pre-post
Hinterberger	2014	49	33	-	45	40-1000 (hrs; range)	Various (inc. MM)	Non pre-post
Huang	2009	23	16	23	31.5	8.4	Zen	Non pre-post
Jo	2014	20	7	19 (from 20)	40.7	3 (minimum)	Zen	Non pre-post
Kasamatsu	1966	48	48	18	24-72 (range)	1-20 (range)	Zen	Non pre-post
Kubota	2001	25	11	-	23.1	0	Zen	Non pre-post
Lahey	2011	18	7	-	18-33 (range)	0	MM	Non pre-post
Lagopoulos	2009	18	13	-	52	9-14 (range)	Acem	Non pre-post
Lehmann	2012	71 (15 Zen)	NR	-	41.4 (Zen=42)	11.3 (Zen=12.3)	Various (inc. Zen)	Non pre-post
Lo	2003	20	NR	10	NR	NR	Zen	Non pre-post
Lo	2013	10	7	10	28	5.8	Zen	Non pre-post
Milz	2014	23	23	2	23.2	0	MM	Non pre-post
Murata	2004	22	22	-	23.3	0	Su-soku	Non pre-post
Pasquini	2015	17	9	14	44.6	2 (minimum)	Zen	Non pre-post
Ren	2011	32	23	16	23.3	0	Su-soku	Non pre-post
Sobolewski	2011	13	7	13	38.7	5 (minimum)	MM	Non pre-post
Stinson	2013	13	NR	-	NR	0	Neurofeedback	Non pre-post
Takahashi	2005	20	20	-	28.6	0	Zen	Non pre-post
Tanaka	2014	10	4	10	49.2	11.6	MM	Non pre-post
Teper	2013	20	9	18	33	3.19	MM	Non pre-post

Running title: The neurophysiology of mindfulness

Teper	2014	45 (from 47)	27	-	19.26	0	Trait mindfulness	Non pre-post
van Leeuwen	2012	8	5	8	29	5	MM	Non pre-post
Yu	2011	15	14	-	38	0	Zen	Non pre-post

Note: MBCT = mindfulness-based cognitive therapy; MM = mindfulness meditation; NCC = neural correlates of consciousness; NR = not recorded; RCT = randomized controlled trial; TM = transcendental meditation; Number of meditators is presented in column headed by Meditators.

Running title: The neurophysiology of mindfulness

Table 2. Demographics of participants with a clinical history

First author	Year	<i>n</i> meditators	<i>n</i> males meditators	<i>n</i> controls	Mean age meditators	Meditation type	Psychiatric Disorder
Barnhofer	2007	10 (from 16)	5	12 (from 18)	48	MBCT	Suicidal depression
Barnhofer	2010	8	1	8	31.6	MM	Previously depressed
Bostanov	2012	32 (from 45)	9	32 (from 46)	50.9	MBCT	Depressed (remission)
Brown	2013	12	NA	15	NA	MM pain manage.	Chronic pain
Garland	2015	11	NA	18	NA	MORE	Chronic pain
Howells	2012	12	2	9	37	MBCT	Bipolar disorder
Keune	2013	40 (from 53)	10	37 (from 50)	48.9	MBCT	Depressed (remission)
Schoenberg (et al.)	2014	26 (from 32)	NA	24 (from 29)	NA	MBCT	ADHD
Schoenberg (& S.)	2014	26	6	25	47.8	MBCT	Depression (current)
Schoenberg (& S.)	2015	26	6	25	47.8	MBCT	Depression (current)

Note: MBCT = mindfulness-based cognitive therapy; MM = mindfulness meditation; MORE = mindfulness-oriented recovery enhancement. All studies featured pre-post designs, and all except Howells were RCTs. All subjects participating had no previous experience with meditation.

Running title: The neurophysiology of mindfulness

Table 3. Results synthesis of papers according to principle findings and according to bandwidth or asymmetry

Outcome	MED > RS	MED = RS	MED < RS	Pre < post (linked to MED)	Pre > post (linked to MED)
Alpha power	Ahani (2014) ; Arita (2012); Cahn (2013); Dunn (1999) ; Hinterberger (2014); Huang (2009); Kasamatsu (1966); Lagopoulos (2009); (2003); Milz (2014) ; Murata (2004; coherence); Takahashi (2005) ; Yu (2011)	Berkovich-Ohana (2013; coherence); Cahn (2010); Kubota (2001) ; Lehmann (2012)	Amihai (2014)		Saggar (2012)
Beta power	Ahani (2014) ; Cahn (2013); Dunn (1999) ; Lo (2003; synchrony)	Cahn (2010); Lagopoulos (2009); Lehmann (2012); Milz (2014) ; Murata (2004; coherence); Yu (2011)	Amihai (2014)	Howells (2012)	Saggar (2012)
Theta power	Ahani (2014) ; Arita (2012); Cahn (2010); Chan (2008); Kasamatsu (1966); Kubota (2001) ; Lagopoulos (2009); Lehmann (2012); Lomas (2014); Takahashi (2005) ; Tanaka (2014)	Amihai (2014); Berkovich-Ohana (2013; coherence); Milz (2014) ; Murata (2004; coherence)	Dunn (1999) ; Huang (2009); Yu (2011)	Howells (2012) ; Xue (2014)	Saggar(2012; in RS); Tang (2009)
Delta power	Cahn (2010; at frontal brain regions)	Amihai (2014); Cahn (2010; at central and parietal brain regions); Lagopoulos (2009); Milz (2014)	Dunn (1999)		
Gamma power	Berkovich-Ohana (2012); Cahn (2010); Hauswald (2015); Lehmann (2012)	Amihai (2014); Berkovich-Ohana (2012; coherence); Milz (2014)			
Greater relative left-sided activation	Amihai (2014); Chan (2008)	Milz et al. (2014)		Barnhofer (2007) ; Barnhofer (2010) ; Davidson (2003)	Keune (2011) ; CNT also decreased)

Note: > = significantly greater than; < = significantly lower than; = = no significant differences; CNT = control group; MED = meditation; RS = resting state. Studies featuring novice participants are indicated by the author/year being italicized in bold. All findings refer to amplitude, unless stated otherwise (e.g., coherence).

Running title: The neurophysiology of mindfulness

Supplementary Table 1. Quality rating of papers with healthy participant samples

First author	Year	Selection bias	Design	Confounders	Blinding	Data collection	Attrition	Global
Ahani	2014	1	1	3	3	1	1	1
Amihai	2014	1	2	2	2	1	2	2
Arita	2012	NA	NA	NA	NA	NA	NA	NA
Becker	1981	1	3	3	3	2	3	3
Berkovich-Ohana	2012	1	1	1	1	1	2	1
Berkovich-Ohana	2013	1	1	1	1	1	2	1
Brown	2010	2	1	3	2	1	2	2
Cahn	2010	1	3	3	3	1	3	3
Cahn	2013	1	3	3	3	1	3	3
Chan	2008	3	3	3	3	1	3	3
Davidson	2003	1	1	1	2	1	1	1
Delgado	2013	2	2	2	3	1	2	2
Dunn	1999	2	3	2	2	3	3	3
Ferrarelli	2013	1	1	1	2	1	2	1
Hauswald	2015	2	2	2	2	1	2	2
Hinterberger	2011	1	2	2	2	1	2	2
Hinterberger	2014	1	2	2	2	1	2	2
Huang	2009	1	1	2	2	1	2	2
Jo	2014	2	1	2	2	1	1	1
Kasamatsu	1966	1	2	2	2	2	2	2
Kerr	2011	1	1	1	2	1	1	1
Kubota	2001	2	2	2	2	1	2	2
Lakey	2011	1	2	2	2	1	2	2
Lagopoulos	2009	1	1	1	2	2	1	1
Lehmann	2012	2	1	2	2	1	2	2
Lo	2003	2	3	3	2	1	2	3
Lo	2013	1	2	2	2	2	2	2
Lomas	2014	1	2	2	2	1	2	2
Milz	2014	1	1	1	2	1	2	1
Moore	2012	1	1	1	1	1	1	1
Murata	2004	1	2	2	2	1	2	2
Pasquini	2015	2	1	2	2	1	2	2
Ren	2011	1	1	2	2	2	2	2
Saggar	2012	1	1	1	2	1	1	1
Slagter	2007	3	3	3	2	1	2	3
Slagter	2009	2	1	1	1	1	2	1
Sobolewski	2011	1	1	1	2	1	2	1
Stinson	2013	2	3	3	1	1	2	3
Takahashi	2005	2	2	2	2	1	2	2
Tanaka	2014	1	1	2	2	1	1	1
Tang	2009	1	1	2	2	1	1	1
Teper	2013	1	1	1	2	1	1	1
Teper	2014	1	2	1	1	1	1	1

Running title: The neurophysiology of mindfulness

van Leeuwen	2012	1	1	1	2	1	2	1
Xue	2014	1	1	2	2	1	2	2
Yu	2011	2	3	2	3	2	2	3

Note: NA = full pdf not available. Studies featuring novice participants are indicated by the author being italicized in bold.

Running title: The neurophysiology of mindfulness

Supplementary Table 2. Quality rating of papers with clinical samples

First author	Year	Selection bias	Design	Confounders	Blinding	Data collection	Attrition	Global
<i>Barnhofer</i>	2007	1	1	2	1	1	1	1
<i>Barnhofer</i>	2010	1	1	1	1	1	1	1
<i>Bostanov</i>	2012	1	2	2	2	1	2	2
<i>Brown</i>	2013	NA	NA	NA	NA	NA	NA	NA
<i>Garland</i>	2015	1	1	1	1	1	2	1
<i>Howells</i>	2012	1	1	1	2	1	2	1
<i>Keune</i>	2013	1	1	1	1	1	1	1
<i>Schoenberg (et al.)</i>	2014	1	1	2	1	1	1	1
<i>Schoenberg (& S.)</i>	2014	1	1	1	1	1	1	1
<i>Schoenberg (& S.)</i>	2015	1	1	1	1	1	1	1

Note: NA = not available. Studies featuring novice participants are indicated by the author being italicized in bold.

Running title: The neurophysiology of mindfulness

Supplementary Table 3. Alpha bandwidth

			Meditation vs resting state	
Author	Year	Meditation	Protocol: Resting	Findings
<i>Higher amplitude during meditation</i>				
Ahani	2014	15 mins MM (E-C)	15 mins audio listening (E-C)	ME > CNT ($F(1,33) = 10.58, p \leq 0.0011$)
Arita	2012	NA	NA	MED > RS (stats NA)
Cahn	2013	21 mins (Vipassana)	21 mins (E-C)	MED > RS ($F(1, 15) = 6.64, p < .05$)
Dunn	1999	15 mins (FA) then 15 mins (MM)	15 mins (E-C)	MED (MM) > RS & FA. Stats NR
Hinterberger	2014	15 mins (self-chosen), 2 mins (MM), 2 mins (thoughtless emptiness), 2 mins (FA on 'third eye'), 2 mins (FA on body axis)	5 mins (E-O), 5 mins (E-C), 5 mins (reading)	MED (MM) > RS ($t = 2.7, p < .05$)
Huang	2009	40 mins (Zen) (vs 40 mins rest for CNT)		EXP (MED) > CNT (rest) ($F(1, 45) = 31.57, P < .0001$)
Kasamatsu	1966	Time NR (Zen)	Time NR (E-C)	MED > RS. Stats NR.
Lagopoulos	2009	20 mins (Acem)	20 mins (E-C)	MED > RS ($F(1, 17) = 7.19, p = .02$)
Lo	2003	40 mins (Zen)	15 mins (E-C)	MED > RS. Stats NR.
Milz	2014	2 x 5 mins (breath counting)	3 x 5 mins (E-C)	Power: MED > RS ($t = 3.02, p = .036$); Coherence: MED = RS ($t(22) = 1.29, p = .20$);
Murata	2004	15 mins (Su-soku; E-O)	15 mins (E-O)	MED > RS (coherence) ($t = 3.03, p < .01$)
Takahashi	2005	15 mins (Su-soku)	15 mins (E-O)	MED > RS ($F(1, 19) = 29.47, p < .001$)
Yu	2011	20 mins (Zen: tanden breathing)	2 mins (E-C)	MED > RS ($F = 9.31, p < .001$)
<i>Lower amplitude during meditation</i>				
Amihai	2014	Therevada = 15 mins (Samatha) & 15 mins (Vipassana); V=Vajrayana = 15 mins (deity) & 15 mins (Rig-pa)	10 mins (E-C)	MED (Therevada) < RS ($F(2,18) = 6.84, p < 0.01$); MED (Vajrayana) = RS ($p > 0.8$)
<i>No significant differences (or no clear reportable patterns)</i>				
Berkovich-Ohana	2013	15 mins (MM)	2.5 mins (E-C) + 2.5 mins (E-O)	MED = RS (coherence); Stats NR. RS: EXP = CNT (coherence); Stats NR.
Cahn	2010	21 mins (Vipassana)	21 mins (E-C)	MED = RS ($F(1, 15) = 0.096, p = .76$)
Hinterberger	2011	15 mins (self-chosen), 2 mins (MM), 2 mins (thoughtless emptiness), 2 mins (FA on 'third eye'), 2 mins (FA on body axis)	5 mins (E-O), 5 mins (E-C), 5 mins (reading)	Complex analyses. Stats NU.
Kubota	2001	25 mins (Su-soku)	2.5 mins (cued-breathing)	MED = RS ($t = 0.68, p$ NR)
Lehmann	2012	60 mins (self-chosen)	4 mins (20 sec E-O, 40 sec E-C; x 4)	MED (Zen) = RS. $T = 1.81, p = 0.09$
Lo	2013	40 mins (Zen)	15 mins (E-C)	Complex analyses. Stats NU
<i>Pre-post changes</i>				
Saggar	2012	12 mins (MM: pre, mid, & post retreat)		Group x time interaction: ($F(2,41) = 23.26, p < .001$): pre-post decrease for EXP ($t(21) = 6.59, p < .001$), not CNT
			Meditation vs task	
Author	Year	Meditation	Protocol: task	Findings
Becker	1981	30 mins (self-chosen MED: Zen, TM or Yoga), then 30 mins (self-chosen MED) concurrent with task	Auditory clicks (30 mins)	Task: EXP = CNT (alpha suppression). Stats NR
Berkovich-Ohana	2013	15 mins (MM)	Time-production task (2-3 mins)	EXP = CNT (coherence); Stats NR
Kerr	2011		Cued attention-detection runs	EXP (vs CNT): Enhanced alpha modulation in early (600-800ms period) (Mann-Whitney, $p < .01$)
Pasquini	2015		Focused attention task	EXP in task: Negative correlation between alpha power and both meditation practice time ($r = -0.52, p = .003$) and meditation weekly frequency ($r = -0.41, p = .021$).

Running title: The neurophysiology of mindfulness

Ren	2011	Time NR (Su-soku)	Problem solving (Time NR)	EXP (MED) < CNT (cognitive task) ($F(2, 45) = 4.14, p = .05$)
Schoenberg (& S.)	2015		Go/NoGo task	Task: pre-post power increase for negative stimuli ($t(24) = 2.58, p = .02$) for CNT only (EXT: no significant increase).
Stinson	2013	Time NR (relaxation – 'Alpha brain state exercise')	Neurofeedback video (Time NR)	MED = task. Stats NR.

Note: > = significantly greater than; < = significantly lower than; = = no significant differences; CNT = control group; E-C = eyes-closed; E-O = eyes-open; EXP = experimental group; FA = focussed-attention (concentrative) meditation; (from ...) = initial number of participants in a pre-post study; MED = meditation; MM = Mindfulness meditation; NA = not available; NR = not reported; NU = not usable (here); NCC = neural correlates of consciousness (i.e., EEG measurement during MED vs RS); ROI = region of interest; RS = resting state. All bandwidth outcomes pertain to power, unless otherwise stated in parentheses (e.g., coherence). Most entries are comparing the experimental group (EXT; i.e., meditators) under different conditions (e.g., RS vs MED): significantly higher power levels during meditation are written as MED > RS; significantly lower levels as MED < RS; and no significant differences as MED = RS. Some entries are comparing two groups (i.e., EXP vs CNT) on a particular condition (e.g., RS): this will be indicated as RS: EXT >/=/< CNT. Studies featuring novice participants are indicated by the author being italicized in bold.

Running title: The neurophysiology of mindfulness

Supplementary Table 4. Beta bandwidth

			Meditation vs resting state	
Author	Year	Meditation	Protocol: Resting	Findings
<i>Higher amplitude during meditation</i>				
Ahani	2014	15 mins MM (E-C)	15 mins audio listening (E-C)	ME > CNT ($F(1,33) = 142.03, p \leq 0.004$)
Cahn	2013	21 mins (Vipassana)	21 mins (E-C)	MED > RS (synchrony) ($F(1, 15) = 9.01, p < .01$)
Dunn	1999	15 mins (FA) then 15 mins (MM)	15 mins (E-C)	MM > RS & FA. Stats NR
Lo	2003	40 mins (Zen)	15 mins (E-C)	MED > RS. Stats NR.
<i>Lower amplitude during meditation</i>				
Amihai	2014	Therevada = 15 mins(Samatha) & 15 mins (Vipassana); V=Vajrayana = 15 mins (deity) & 15 mins (Rig-pa)	10 mins (E-C)	MED (Therevada) < RS ($F(2,18) = 3.68, p < 0.05$); MED (Vajrayana) < RS ($F(2,18) = 8.42, p < 0.01$)
<i>No significant differences (or no clear reportable patterns)</i>				
Cahn	2010	21 mins (Vipassana)	21 mins (E-C)	MED = RS ($F(1, 15) = 0.62, p = .44$)
Hinterberger	2011	15 mins (self-chosen), 2 mins (MM), 2 mins (thoughtless emptiness), 2 mins (FA on 'third eye'), 2 mins (FA on body axis)	5 mins (E-O), 5 mins (E-C), 5 mins (reading)	Complex analyses. Stats NU.
Lagopoulos	2009	20 mins (Acem)	20 mins (E-C)	MED = RS ($F(1, 17) = 0.57, p = .46$)
Lehmann	2012	60 mins (self-chosen)	4 mins (20 sec E-O, 40 sec E-C; x 4)	MED (Zen) = RS. $T = 0.48, p = 0.63$
Milz	2014	2 x 5 mins (breath counting)	3 x 5 mins (E-C)	Power: MED = RS; Stats NR. Coherence: MED = RS ($t(22) = 0.11, p = .91$)
Murata	2004	15 mins (Su-soku; E-O)	15 mins (E-O)	EXP: MED = RS (coherence). Stats NR.
Yu	2011	20 mins (Zen: tanden breathing)	2 mins (E-C)	MED = RS ($F = 0.96, p = .44$)
<i>Pre-post changes</i>				
Saggar	2012	12 mins (MM: pre, mid, & post retreat)		Group x time interaction: ($F(2,41) = 7.11, p < .01$): pre-post decrease for EXP ($t(21) = 8.65, p < .001$), not CNT
Howells	2012		3 mins (E-O), 3 mins (E-C),	RS (EXP only): post < pre ($t = 2.23, p < .05$)
			Meditation vs task	
Author	Year	Meditation	Protocol: task	Findings
Stinson	2013	Time NR (relaxation – 'Alpha brain state exercise')	Neurofeedback video (Time NR)	MED < task. Stats NR.

Note: > = significantly greater than; < = significantly lower than; = = no significant differences; CNT = control group; E-C = eyes-closed; E-O = eyes-open; EXP = experimental group; FA = focussed-attention (concentrative) meditation; (from ...) = initial number of participants in a pre-post study; MED = meditation; MM = Mindfulness meditation; NA = not available; NR = not reported; NU = not usable (here); NCC = neural correlates of consciousness (i.e., EEG measurement during MED vs RS); ROI = region of interest; RS = resting state. All bandwidth outcomes pertain to power, unless otherwise stated in parentheses (e.g., coherence). Most entries are comparing the experimental group (EXT; i.e., meditators) under different conditions (e.g., RS vs MED): significantly higher power levels during meditation are written as MED > RS; significantly lower levels as MED < RS; and no significant differences as MED = RS. Some entries are comparing two groups (i.e., EXP vs CNT) on a particular condition (e.g., RS): this will be indicated as RS: EXT >/=< CNT. Studies featuring novice participants are indicated by the author being italicized in bold.

Running title: The neurophysiology of mindfulness

Supplementary Table 5. Theta bandwidth

			Meditation vs resting state	
Author	Year	Meditation	Protocol: Resting	Findings
<i>Higher amplitude during meditation</i>				
Ahani	2014	15 mins MM (E-C)	15 mins audio listening (E-C)	ME > CNT ($F(1,33) = 118.79, p \leq 0.001$)
Arita	2012	NA	NA	MED > RS (stats NA)
Cahn	2010	21 mins (Vipassana)	21 mins (E-C)	MED > RS (condition x ROI interaction – only at specific sites ($F(2, 30) = 7.75, p = .002$). Increase in MED at Fz ($p = .006$), but not Cz ($p = .99$) or Pz ($p = .85$))
Chan	2008	12 mins (Triarchic Body Relaxation Technique)	5 mins (E-C)	MED > RS (a range of t-tests; $t = -3.73 - -4.82, p < .02$).
Kasamatsu	1966	Time NR (Zen)	Time NR (E-C)	MED > RS. Stats NR.
Kubota	2001	25 mins (Su-soku)	2.5 mins (cued-breathing)	MED > RS ($t = 6.14, p < .0001$)
Lagopoulos	2009	20 mins (Acem)	20 mins (E-C)	MED > RS ($F(1, 17) = 4.99, p = .04$)
Lehmann	2012	60 mins (self-chosen)	4 mins (20 sec E-O, 40 sec E-C; x 4)	MED (Zen) > RS. $T = 4.95, p < 0.001$.
Takahashi	2005	15 mins (Su-soku)	15 mins (E-O)	MED > RS ($F(1, 19) = 5.5, p = .031$)
Tanaka	2014	40 mins (MM)	8 mins (E-C)	EXP: MED > RS (stats NA). MED: EXP > CNT ($p < .0001$). RS: EXP < CNT ($p < .0001$).
<i>Lower amplitude during meditation</i>				
Dunn	1999	15 mins (FA) then 15 mins (MM)	15 mins (E-C)	MM < RS; Stats NR; MM > FA; Stats NR
Huang	2009	40 mins (Zen) (vs 40 mins rest for CNT)		EXP (MED) < CNT (rest) ($F(1, 45) = 28.68, P < .0001$)
Yu	2011	20 mins (Zen: tanden breathing)	2 mins (E-C)	MED < RS ($F = 9.85, p < .001$)
<i>No significant differences (or no clear reportable patterns)</i>				
Amihai	2014	Therevada = 15 mins (Samatha) & 15 mins (Vipassana); Vajrayana = 15 mins (deity) & 15 mins (Rig-pa)	10 mins (E-C)	MED (Therevada) = RS ($F(2,18) = 1.11, p > 0.3$); MED (Vajrayana) = RS ($F(2,16) = 2.5, p > 0.1$)
Berkovich-Ohana	2013	15 mins (MM)	2.5 mins (E-C) + 2.5 mins (E-O)	MED = RS (coherence); Stats NR. RS: EXP = CNT (coherence); Stats NR.
Hinterberger	2011	15 mins (self-chosen), 2 mins (MM), 2 mins (thoughtless emptiness), 2 mins (FA on 'third eye'), 2 mins (FA on body axis)	5 mins (E-O), 5 mins (E-C), 5 mins (reading)	Complex analyses. Stats NU.
Milz	2014	2 x 5 mins (breath counting)	3 x 5 mins (E-C)	Power: MED = RS; Stats NR. Coherence: MED = RS ($t(22) = 0.62, p = .53$)
Murata	2004	15 mins (Su-soku; E-O)	15 mins (E-O)	EXP: MED = RS (coherence). Stats NR.
<i>Pre-post changes</i>				
Lomas	2014	10 mins (MM)	5 mins (E-C)	Pre: MED > RS ($F(1, 27) = 7.14, p = .013$); Post: MED > RS ($F(1, 27) = 5.74, p = .024$)
Tang	2009		Time NR (E-C)	Group x time interaction ($F(1, 32) = 4.92, p < .05$): pre-post decrease for EXP ($p < .05$), not CNT
Xue	2014		5 mins (E-C)	Group x time interaction (in connectivity) ($F(1,43) = 2.93; p = 0.09$). Pre: EXP = CNT ($p > .05$). Post: decreased path-length in EXP ($t(23) = 3.72, p = .001$), not CNT.
			Meditation vs task	
Author	Year	Meditation	Protocol: task	Findings
Berkovich-Ohana	2013	15 mins (MM)	Time-production task (2-3 mins)	EXP = CNT (coherence); Stats NR
Howells	2012	Resting state: 3 mins (E-O), 3 mins (E-C)	Sustained attention (visual A-	RS (EXP only): post < pre ($t = 2.29, p < .05$)

Running title: The neurophysiology of mindfulness

			X continuous performance) (10 mins)	
Pasquini	2015		Focused attention task	EXP in task: Correlation between theta power and meditation weekly frequency ($r = 0.42, p = .02$).
Slagter	2009		Attentional blink (time NR)	Group x time interaction: MED-related changes in the phase of target induced EEG responses. Stats NU
Stinson	2013	Time NR (relaxation – 'Alpha brain state exercise')	Watching video – explaining neurofeedback (Time NR)	MED < task. Stats NR.

Note: > = significantly greater than; < = significantly lower than; = = no significant differences; CNT = control group; E-C = eyes-closed; E-O = eyes-open; EXP = experimental group; FA = focussed-attention (concentrative) meditation; (from ...) = initial number of participants in a pre-post study; MED = meditation; MM = Mindfulness meditation; NA = not available; NR = not reported; NU = not usable (here); NCC = neural correlates of consciousness (i.e., EEG measurement during MED vs RS); ROI = region of interest; RS = resting state. All bandwidth outcomes pertain to power, unless otherwise stated in parentheses (e.g., coherence). Most entries are comparing the experimental group (EXT; i.e., meditators) under different conditions (e.g., RS vs MED): significantly higher power levels during meditation are written as MED > RS; significantly lower levels as MED < RS; and no significant differences as MED = RS. Some entries are comparing two groups (i.e., EXP vs CNT) on a particular condition (e.g., RS); this will be indicated as RS: EXT >/=/< CNT. Studies featuring novice participants are indicated by the author being italicized in bold.

Running title: The neurophysiology of mindfulness

Supplementary Table 6. Delta bandwidth

			<i>Meditation vs resting state</i>	
Author	Year	Meditation	Protocol: Resting	Findings
<i>Higher amplitude during meditation</i>				
Cahn	2013	21 mins (Vipassana)	21 mins (E-C)	Expertise x state x ROI interaction(synchrony) ($F(2, 2, 28) = 5.83, p < .01$): long- term EXP higher synchrony during MED (vs RS) at frontal sites, but not central or parietal sites; for short term meds, MED = RS
<i>Lower amplitude during meditation</i>				
Dunn	1999	15 mins (FA) then 15 mins (MM)	15 mins (E-C)	MM < RS; Stats NR
<i>No significant differences (or no clear reportable patterns)</i>				
Amihai	2014	Therevada = 15 mins (Samatha) & 15 mins (Vipassana); Vajrayana = 15 mins (deity) & 15 mins (Rig-pa)	10 mins (E-C)	Med < RS ($F(2,18) = 8.37, p < 0.01$). Post-hoc: no diff between– RS and Therevada MED (only Vajrayana MED)
Cahn	2010	21 mins (Vipassana)	21 mins (E-C)	MED = RS ($F(1, 15) = 1.85, p = .19$)
Hinterberger	2011	15 mins (self-chosen), 2 mins (MM), 2 mins (thoughtless emptiness), 2 mins (FA on 'third eye'), 2 mins (FA on body axis)	5 mins (E-O), 5 mins (E-C), 5 mins (reading)	Complex analyses. Stats NU.
Lagopoulos	2009	20 mins (Acem)	20 mins (E-C)	MED = RS ($F(1, 17) = 0.99, p = .34$)
Milz	2014	2 x 5 mins (breath counting)	3 x 5 mins (E-C)	Power: MED = RS. Stats NR
			<i>Meditation vs task</i>	
Author	Year	Meditation	Protocol: task	Findings
Stinson	2013	Time NR (relaxation – 'Alpha brain state exercise')	Watching video – explaining neurofeedback (Time NR)	MED < task. Stats NR.

Note: > = significantly greater than; < = significantly lower than; = = no significant differences; CNT = control group; E-C = eyes-closed; E-O = eyes-open; EXP = experimental group; FA = focussed-attention (concentrative) meditation; (from ...) = initial number of participants in a pre-post study; MED = meditation; MM = Mindfulness meditation; NA = not available; NR = not reported; NU = not usable (here); NCC = neural correlates of consciousness (i.e., EEG measurement during MED vs RS); ROI = region of interest; RS = resting state. All bandwidth outcomes pertain to power, unless otherwise stated in parentheses (e.g., coherence). Most entries are comparing the experimental group (EXT; i.e., meditators) under different conditions (e.g., RS vs MED): significantly higher power levels during meditation are written as MED > RS; significantly lower levels as MED < RS; and no significant differences as MED = RS. Some entries are comparing two groups (i.e., EXP vs CNT) on a particular condition (e.g., RS): this will be indicated as RS: EXT >/=/
 CNT. Studies featuring novice participants are indicated by the author being italicized in bold.

Running title: The neurophysiology of mindfulness

Supplementary Table 7. Gamma bandwidth

			<i>Meditation vs resting state</i>	
Author	Year	Meditation	Protocol: Resting	Findings
<i>Higher amplitude during meditation</i>				
Berkovich-Ohana	2012	15 mins (MM)	2.5 mins (E-C) + 2.5 mins (E-O)	MED > RS ($F(1, 34) = 17.00, p < 0.0001$).
Cahn	2010	21 mins (Vipassana)	21 mins (E-C)	MED > RS ($F(1, 15) = 9.32, p = .008$).
Hauswald	2015	20 mins (Zen, E-O)	5 mins (E-O)	Increased gamma power in MED correlated both with trait mindfulness ($p = .015$) and years of practice ($p = .036$).
Lehmann	2012	60 mins (self-chosen)	4 mins (20 sec E-O, 40 sec E-C; x 4)	MED (Zen) > RS. $T = 2.66, p = 0.019$.
<i>No significant differences (or no clear reportable patterns)</i>				
Amihai	2014	Therevada = 15 mins (Samatha) & 15 mins (Vipassana); Vajrayana = 15 mins (deity) & 15 mins (Rig-pa)	10 mins (E-C)	MED (Therevada) = RS; Stats NR. MED (Vajrayana) < RS ($F(2,16) = 6.16, p > 0.01$)
Berkovich-Ohana	2013	15 mins (MM)	2.5 mins (E-C) + 2.5 mins (E-O)	MED = RS (coherence); Stats NR. RS: EXP = CNT (coherence); Stats NR.
Hinterberger	2011	15 mins (self-chosen), 2 mins (MM), 2 mins (thoughtless emptiness), 2 mins (FA on 'third eye'), 2 mins (FA on body axis)	5 mins (E-O), 5 mins (E-C), 5 mins (reading)	Complex analyses. Stats NU.
Milz	2014	2 x 5 mins (breath counting)	3 x 5 mins (E-C)	Power; MED = RS. Stats NR
			<i>Meditation vs task</i>	
Author	Year	Meditation	Protocol: task	Findings
Berkovich-Ohana	2013	15 mins (MM)	Time-production task (2-3 mins)	EXP = CNT (coherence); Stats NR
Ferarelli	2013		Sleep	Correlation: MED experience & NREM gamma ($r = 0.47, p = .017$). No correlation with REM gamma
Schoenberg (& S.)	2015		Go/NoGo task	Task: Interaction (time x group x site x epoch) ($F(1, 47) = 4.12, p = .05$) – pre-post power increase for CNT only (EXT: no increase).
Stinson	2013	Time NR (relaxation – 'Alpha brain state exercise')	Watching video – explaining neurofeedback (Time NR)	MED < task. Stats NR.

Note: > = significantly greater than; < = significantly lower than; = = no significant differences; CNT = control group; E-C = eyes-closed; E-O = eyes-open; EXP = experimental group; FA = focussed-attention (concentrative) meditation; (from ...) = initial number of participants in a pre-post study; MED = meditation; MM = Mindfulness meditation; NA = not available; NR = not reported; NU = not usable (here); NCC = neural correlates of consciousness (i.e., EEG measurement during MED vs RS); ROI = region of interest; RS = resting state. All bandwidth outcomes pertain to power, unless otherwise stated in parentheses (e.g., coherence). Most entries are comparing the experimental group (EXT; i.e., meditators) under different conditions (e.g., RS vs MED); significantly higher power levels during meditation are written as MED > RS; significantly lower levels as MED < RS; and no significant differences as MED = RS. Some entries are comparing two groups (i.e., EXP vs CNT) on a particular condition (e.g., RS): this will be indicated as RS: EXT >/=< CNT. Studies featuring novice participants are indicated by the author being italicized in bold.

Running title: The neurophysiology of mindfulness

Supplementary Table 8. Asymmetry Findings

Author	Year	Meditation	Meditation vs resting state	
			Protocol	Findings
<i>Increase in relative left-frontal activation linked to meditation</i>				
Amihai	2014	Therevada = 15 mins (Samatha) & 15 mins (Vipassana); Vajrayana = 15 mins (deity) & 15 mins (Rig-pa)	10 mins resting state (E-C)	MED (Therevada): Condition x location interaction ($F(4,36) = 3.09, p < 0.05$). MED (Vipassana) > RS at left location ($p < 0.05$), but not right or center ($p > 0.2$)
Barnhofer	2007	8 x 1min (4 = E-O, 4 = E-C)	8 x 1min (4 = E-O, 4 = E-C)	Group x time interaction ($F(2, 19) = 3.7, p = .044$): pre-post decreases in relative left prefrontal asymmetry for CNT ($p = .003$), but not EXP ($p = .918$)
Barnhofer	2010	EXP = 15 mins (MM); CNT = 15 mins (LKM)	2 mins (E-C): EEG assessed pre and post MED	Pre-post increase (EXP & CNT) in relative left prefrontal activation ($F(1, 13) = 5.06, p = .04$)
Chan	2008	Triarchic Body Relaxation Technique (12 mins) 5 mins (E-C)	5 mins resting state (E-C)	MED > RS (left-sided activation); $F(1, 18) = 5.42, p = .032$
Davidson	2003	8 x 1min (4 = E-O, 4 = E-C)	Writing about experiences (EEG recorded 1 min before & 3 mins after)	Group x time interaction in RS ($F(1, 37) = 5.14, p < .05$): pre-post increase in relative left- activation for EXP (not CNT)
<i>Decrease in relative left-frontal activation linked to meditation</i>				
Keune	2011		Sad mood induction (sad music, and neg. experience recall). Time NR.	Pre-post decrease (EXP & CNT) in relative left prefrontal activation ($F(4, 64) = 3.38, p < .05$)
<i>No change in relative left-frontal activation linked to meditation</i>				
Milz	2014	2 x 5 mins (breath counting)	3 x 5 mins resting state (E-C)	Power: MED = RS. Stats NR

Note: > = significantly greater than; < = significantly lower than; = = no significant differences; CNT = control group; E-C = eyes-closed; E-O = eyes-open; EXP = experimental group; FA = focussed-attention (concentrative) meditation; (from ...) = initial number of participants in a pre-post study; MED = meditation; MM = Mindfulness meditation; NA = not available; NR = not reported; NU = not usable (here); NCC = neural correlates of consciousness (i.e., EEG measurement during MED vs RS); ROI = region of interest; RS = resting state. All bandwidth outcomes pertain to power, unless otherwise stated in parentheses (e.g., coherence). Most entries are comparing the experimental group (EXT; i.e., meditators) under different conditions (e.g., RS vs MED): significantly higher power levels during meditation are written as MED > RS; significantly lower levels as MED < RS; and no significant differences as MED = RS. Some entries are comparing two groups (i.e., EXP vs CNT) on a particular condition (e.g., RS): this will be indicated as RS: EXT >/=< CNT. Studies featuring novice participants are indicated by the author being italicized in bold.

Running title: The neurophysiology of mindfulness

Supplementary Table 9. ERP Findings

Author	Year	Meditation	Meditation vs task	
			Protocol: task	Findings
Bostanov	2012	20 mins (MM)	Auditory stimuli during MM	Increased pre-post 'Late contingent negative variation for EXP. but not CNT ($F = 7.7, p < .01$)
Brown	2010		Stimulation of pain (5 mins)	Lower activation for EXP than CNT in S2 and insula during pain stimulus, $t = 2.51, p < .05$
Brown	2013	-	Anticipation and simulation of pain (time NR)	Anticipatory and pain-evoked ERPs to acute pain reduced in EXP but not CNT (stats NA)
Delgado	2013		Auditory oddball task	Two-way P3b interaction (task \times oddball order): significant effects of meditation after the meditation/control task ($p = 0.01$).
Garland	2015		Event-related affective picture viewing task	Tim \times group \times cue interaction ($F(1, 25) = 4.99, p = .035$). EXP group (vs CNT) = pre-post increases in LPP activation to natural reward cues across 400 – 1000 ms window.
Jo	2014		Performing voluntary finger movement (time NR)	MED = CNT (Readiness Potential amplitude prior to voluntary action) ($p = .26$).
Lahey	2011	EXP = 6 mins (MM); CNT = 6 mins (non-MM-task)	P300-based brain-computer interface (BCI) task	Task: EXP > CNT (P300 amplitude peaks) ($t(16) = 2.10, P < .05$)
Moore	2012		Stroop (time NR)	Group \times time interaction: pre-post increase in focusing attention (EXP only). Stats NU.
Schoenberg (et al.)	2014		Go/NoGo task	Time \times condition \times group interaction: significant pre-post increases for EXP in Go-P3 ($t(23) = -2.986, p = .007$) and NoGo-P3 ($t(23) = -2.502, p = .02$) amplitude at Pz, contrary to pre-post parietal decreases for CNT in Go-P3 ($p = .42$) and NoGo-P3 ($p = .40$).
Schoenberg (& S.)	2014		Go/NoGo task	Pre-post increase in event-related theta synchronization during the late time window (400-800 ms) for EXT ($F(1, 49) = 10.933, p = .002$), vs pre-post decrease for CNT
Slagter	2007		Attentional blink (time NR)	Group \times time interaction ($F(1, 20) = 5.4, p = .03$): pre-post decrease in elicited P3B amplitude for EXP, not CNT
Sobolewski	2011		Looking at emotional pictures (Time NR)	Group \times valence interaction ($p = .03$): EXP less affected by negative emotional load.
Teper	2013		Stroop (time NR)	Task: EXP > CMT (higher amplitude error-related negativity) ($F(1, 36) = 3.32, p < .04$)
Teper	2014		Performance feedback (neutral, aversive, and rewarding) (time NR)	Trait MM: predicts less differentiation of rewarding from neutral feedback. Stats NU
Van Leeuwen	2012		Target detection (time NR)	Task: MED > CNT (enhanced attentional processing). Stats NU.

Note: > = significantly greater than; < = significantly lower than; = = no significant differences; CNT = control group; E-C = eyes-closed; E-O = eyes-open; EXP = experimental group; FA = focussed-attention (concentrative) meditation; (from ...) = initial number of participants in a pre-post study; MED = meditation; MM = Mindfulness meditation; NA = not available; NR = not reported; NU = not usable (here); NCC = neural correlates of consciousness (i.e., EEG measurement during MED vs RS); ROI = region of interest; RS = resting state. All bandwidth outcomes pertain to power, unless otherwise stated in parentheses (e.g., coherence). Most entries are comparing the experimental group (EXT; i.e., meditators) under different conditions (e.g., RS vs MED): significantly higher power levels during meditation are written as MED > RS; significantly lower levels as MED < RS; and no significant differences as MED = RS. Some entries are comparing two groups (i.e., EXP vs CNT) on a particular condition (e.g., RS): this will be indicated as RS: EXT >/=< CNT. Studies featuring novice participants are indicated by the author being italicized in bold.